



HAL
open science

Temperature distribution in Mars atmosphere as measured by the ACS-MIR/ExoMars-TGO solar occultations

Denis Belyaev, Anna Fedorova, Alexander Trokhimovskiy, Oleg Korablev, Franck Montmessin, Kevin Olsen, Juan Alday, Jean-Loup Bertaux

► **To cite this version:**

Denis Belyaev, Anna Fedorova, Alexander Trokhimovskiy, Oleg Korablev, Franck Montmessin, et al.. Temperature distribution in Mars atmosphere as measured by the ACS-MIR/ExoMars-TGO solar occultations. EPSC-DPS Joint Meeting 2019, Sep 2019, Geneva, Switzerland. pp.EPSC-DPS2019-1566-1. insu-04450667

HAL Id: insu-04450667

<https://insu.hal.science/insu-04450667v1>

Submitted on 10 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Temperature distribution in Mars atmosphere as measured by the ACS-MIR/ExoMars-TGO solar occultations

Denis Belyaev (1), Anna Fedorova (1), Alexander Trokhimovskiy (1), Oleg Korablev (1), Franck Montmessin (2), Kevin Olsen (2), Juan Alday (3), Jean-Loup Bertaux (2) and the ACS Science Team

(1) Space Research Institute (IKI), Moscow, Russia, (2) LATMOS/CNRS, Paris, France, (3) AOPP, Department of Physics, University of Oxford, UK, (bdenya.iki@gmail.com)

Abstract

We report results from the first year of temperature distribution retrievals in Mars atmosphere from the solar occultation measurements, being performed by the Atmospheric Chemistry Suite (ACS) onboard the ExoMars Trace Gas Orbiter (TGO). Statistics of observations provides almost regular time coverage including the beginning of the Martian global dust storm in June 2018 ($L_S \approx 190$, MY34). Temperature variability before and during the storm is studied at altitudes from 3-30 km up to 80-100 km.

1. Introduction

The ACS instrument began nominal science operations in April 2018 on board the TGO of the ExoMars mission. The mid-infrared channel (MIR) of ACS is a cross-dispersion echelle spectrometer dedicated to solar occultation measurements in the 2.3–4.2 μm range [1]. This experiment achieves the signal-to-noise ratio $\text{SNR} \sim 3000$ with the instrumental resolving power of $>30,000$. It is able to accomplish the most sensitive measurements of the trace gases ever present in the Martian atmosphere. In parallel, the tiniest atmospheric layers – even up to 200 km – may be probed along the occultation line of sight measuring in strong CO_2 absorption bands (e.g. 2.7 μm , and 4.3 μm).

Each occultation session covers a spectral interval with one or a few CO_2 absorption bands appropriate for density, scale height and temperature retrievals in the Martian atmosphere. In the current study we use several set of spectroscopic lines (e.g. at 2.5–2.6 μm , $\sim 3.15 \mu\text{m}$, and $\sim 4 \mu\text{m}$) and the hydrostatic equilibrium condition to determine temperature profiles from the retrieved CO_2 local densities.

At the moment, we present results from the first year of ACS MIR temperature profile retrievals in the solar occultation mode. Statistics of observations provides almost regular time coverage including the beginning of the Martian global dust storm in June 2018 ($L_S \approx 190$, MY34). Temperature variability before and during the storm is studied at altitudes from 3-30 km up to 80-100 km.

2. Measurements

ACS Middle InfraRed channel (ACS-MIR) is a cross-dispersion echelle spectrometer, operating in the 2.3–4.2 μm range [1]. The echelle orders are dispersed along the x-axis and separated along the y-axis of the focal plane by a secondary grating, making full use of the 2-D detector array.

The spectrometer is devoted to solar occultation measurements to perform vertical profiling of the atmosphere with altitude field of view around 0.5–2 km. Each occultation occurs at a fixed position of the secondary grating that cover spectral interval of $\sim 0.3 \mu\text{m}$ including 10–15 diffraction orders of the echelle grating. Figure 1 demonstrates an example of CO_2 transmission spectra measured and best fitted in two MIR positions: 5 at 2.51 μm and 12 at 3.15 μm .



Figure 1: Examples of ACS-MIR spectral windows in diffraction orders 237 at 2.51 μm (left), and 189–190 at 3.15 μm (right), measured at tangent heights 12 and 19 km, used for the retrieval of CO_2 density.

3. Conclusions

Retrievals of CO₂ density are possible from many spectral windows (echelle orders) of ACS-MIR, and in parallel with minor species. Temperature is extracted from the hydrostatic equilibrium condition, and it gives us link to the dust storm behaviour with altitude, temporal and latitude distributions.

Validation of our results with analogous studies, e.g. MCS/MRO [2] and NOMAD/TGO [3], are in progress.

Acknowledgements

ExoMars is the space mission of ESA and Roscosmos. The ACS experiment is led by IKI Space Research Institute in Moscow. The project acknowledges funding by Roscosmos and CNES. Science operations of ACS are funded by Roscosmos and ESA. Science support in IKI is funded by the Russian Government Grant ##14.W03.31.0017.

References

- [1] Korabiev O. et al. The Atmospheric Chemistry Suite (ACS) of Three Spectrometers for the ExoMars 2016 Trace Gas Orbiter. *Space Sci. Rev.*, 214:7, 2018. <https://doi.org/10.1007/s11214-017-0437-6>

- [2] McCleese D.J. et. Al. Mars Climate Sounder: An investigation of thermal and water vapor structure, dust and condensate distributions in the atmosphere, and energy balance of the polar regions. *Journal of Geophysical Research*, Volume 112, Issue E5, CiteID E05S06, 2007.

- [3] Vandaele A.C. et al. NOMAD, an Integrated Suite of Three Spectrometers for the ExoMars Trace Gas Mission: Technical Description, Science Objectives and Expected Performance. *Space Sci Rev.*, 214:80, 2018. <https://doi.org/10.1007/s11214-018-0517-2>