



**HAL**  
open science

## Simulation of the Hydrogen escape from Mars using a Global Climate Model

Francisco González-Galindo, Jean-Yves Chaufray, Gabriella Gilli, Margaux Vals, Franck Lefèvre, Franck Montmessin, Loïc Rossi, Francois Forget, Ehouarn Millour, Miguel Ángel López-Valverde, et al.

► **To cite this version:**

Francisco González-Galindo, Jean-Yves Chaufray, Gabriella Gilli, Margaux Vals, Franck Lefèvre, et al.. Simulation of the Hydrogen escape from Mars using a Global Climate Model. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. pp.EPSC2022-693, 10.5194/epsc2022-693 . insu-04451192

**HAL Id: insu-04451192**

**<https://insu.hal.science/insu-04451192>**

Submitted on 11 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Europlanet Science Congress 2022  
Palacio de Congresos de Granada, Spain  
18 September – 23 September 2022

EPSC Abstracts

Vol. 16, EPSC2022-693, 2022, updated on 09 Jan 2024

<https://doi.org/10.5194/epsc2022-693>

Europlanet Science Congress 2022

© Author(s) 2024. This work is distributed under  
the Creative Commons Attribution 4.0 License.



# Simulation of the Hydrogen escape from Mars using a Global Climate Model

Francisco González-Galindo et al.

## Introduction

The thermal (Jeans) escape of Hydrogen accumulated during the history of Mars has been one of the major mechanisms explaining the transition of Mars from a thicker and wetter atmosphere in the past to the current thin and dry atmosphere (Brain et al., 2017). Recent observations (Heavens et al., 2018, Chaffin et al., 2021) have revealed a clear link between the water cycle in the lower atmosphere, the transport of water to the middle/upper atmosphere, and the thermal escape of Hydrogen. However, many unknowns remain, including the role of the different processes responsible of transporting water from the lower to the upper atmosphere and converting it to Hydrogen atoms, or the effects of global dust storms (GDS hereafter) compared to the regular seasonal variability.

While different 1D models have been used to reproduce and understand some of the observations (e.g. Chaffin et al., 2017), until now global models have failed to reproduce the observed variability of the H escape. In particular, a recent study with the Laboratoire de Météorologie Dynamique Mars Global Climate Model (LMD-MGCM hereafter) evidences that the model significantly underestimates the H escape rate when comparing with Mars Express SPICAM observations, in particular during the perihelion season (Chaufray et al., 2021).

In this work we will summarize the recent improvements that we have included in the LMD-MGCM in order to better reproduce the observed Hydrogen escape rate, and will discuss some of the results obtained with the improved model.

## Model description

We have included three improvements with respect to the version of the LMD-MGCM used in Chaufray et al., 2021.

First, we have incorporated in the simulations a sophisticated model of the microphysics of water ice clouds allowing for the formation of supersaturated water layers (Navarro et al., 2014). Second, we have extended the photochemical model in the LMD-MGCM to incorporate the chemistry of H<sub>2</sub>O<sup>+</sup> and derived ions, as well as of deuterated (both neutral and ion) species. Third, we have also included in the calculations an improved model of deuterium fractionation (Vals et al., 2022). While this allows us to study the D escape, we will focus here only on the H escape; simulations of the deuterium escape are discussed in Chaufray et al. (this issue).

## Preliminary results

The incorporation in the calculations of the microphysical model allowing for the formation of supersaturated water layers significantly increases the amount of water in the upper atmosphere of the planet with respect to the previous calculations, producing a strong enhancement of up to one order of magnitude in the H escape rate. The incorporation of the chemistry of water-derived ions further increases the escape rate in between ~20 and ~40%, depending on the season. This results in a better agreement with observations of H escape (figure 1). However, significant differences still remain. In particular, the decrease in the rate of H escape at the end of the year is not well captured by the model, suggesting that, in the model, water remains in the upper atmosphere longer than observed.

We study also the interannual variability of the simulated escape rate. While the solar activity seems to play a secondary role, dust storms in the lower atmosphere have a clear effect over the H escape rate. Our simulations show, for example, that the global dust storm in MY34 increased the annually integrated H escape rate in about 30%. This confirms the importance of taking into account the effects of GDSs when calculating the accumulated escape rate over Martian history.

This work opens the doors to studying the H escape rate at past Mars conditions characterized by different orbital parameters (e.g. obliquity, time of perihelion, etc.). See Gilli et al., this issue, for a first study in this direction.




Figure 1.  $H$  escape rate simulated for MY28 (light green line) and MY33 (orange line). The black thin line shows the escape rate for MY28 simulated with the previous model version, taken from Chaufray et al. (2021). The green and red symbols represent measured values of the  $H$  escape rate during MY28 and MY33, respectively, taken from Chaffin et al. (2014) and Heavens et al. (2018)

### Acknowledgements

F.G.-G. and G.G. are funded by the Spanish Ministerio de Ciencia, Innovación y Universidades, the Agencia Estatal de Investigación and EC FEDER funds under project RTI2018-100920-J-I00. AB and MALV were supported by grant PGC2018-101836-B-100 (MCIU/AEI/FEDER, EU). The IAA team acknowledges financial support from the State Agency for Research of the Spanish MCIU through the Center of Excellence "Severo Ochoa" award to the Instituto de Astrofísica de Andalucía (SEV-2017-0709)

### References

- Brain, D., et al., (2017), Solar Wind Interaction and Atmospheric Escape. Chapter 15 in "The Atmosphere and Climate of Mars", Cambridge University Press, doi:10.1017/9781139060172.015
- Chaffin, M., et al. (2021), Martian water loss to space enhanced by regional dust storms, Nat. Astron. doi:10.1038/s41550-021-01425-w
- Chaufray, J.-Y., et al. (2021), Study of the hydrogen escape rate at Mars during martian years 28 and 29 from comparisons between SPICAM/Mars express observations and GCM-LMD simulations. Icarus, doi:10.1016/j.icarus.2019.113498
- Heavens, N., et al. (2018). Hydrogen escape from Mars enhanced by deep convection in dust storms. Nat. Astron., doi:10.1038/s41550-017-0353-4
- Navarro, T., et al. (2014), Global climate modeling of the Martian water cycle with improved microphysics and radiatively active water ice clouds. JGR (Planets), doi.org:10.1002/2013JE004550
- Vals, M., et al. (2022), Improved modeling of Mars' H<sub>2</sub>O cycle using a Mars' Global Climate Model. Paper submitted to JGR-Planets.

**How to cite:** González-Galindo, F., Chaufray, J.-Y., Gilli, G., Vals, M., Lefèvre, F., Montmessin, F., Rossi, L., Forget, F., Millour, E., López-Valverde, M. Á., and Brines, A.: Simulation of the Hydrogen escape from Mars using a Global Climate Model, Europlanet Science Congress 2022, Granada, Spain, 18–23 Sep 2022, EPSC2022-693, <https://doi.org/10.5194/epsc2022-693>, 2022.



## Discussion

We are sorry, but the discussion is only available for users who registered for the conference. Thank you.