



**HAL**  
open science

# The Venus Emissivity Mapper - Investigating the Atmospheric Structure and Dynamics of Venus' Polar Region

Thomas Widemann, Emmanuel Marcq, C. Tsang, N. Mueller, D. Kappel, J. Helbert, M. D. Dyar

► **To cite this version:**

Thomas Widemann, Emmanuel Marcq, C. Tsang, N. Mueller, D. Kappel, et al.. The Venus Emissivity Mapper - Investigating the Atmospheric Structure and Dynamics of Venus' Polar Region. 49th Lunar and Planetary Science Conference, Mar 2018, The Woodlands, United States. pp.LPI Contribution No. 2083, id.2386. insu-04452860

**HAL Id: insu-04452860**

**<https://insu.hal.science/insu-04452860>**

Submitted on 12 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**THE VENUS EMISSIVITY MAPPER – INVESTIGATING THE ATMOSPHERIC STRUCTURE AND DYNAMICS OF VENUS' POLAR REGION.** T. Widemann<sup>1,2</sup>, E. Marcq<sup>3,2</sup>, C. Tsang<sup>4</sup>, N. Mueller<sup>5</sup>, D. Kappel<sup>6</sup>, J. Helbert<sup>6</sup>, M. D. Dyar<sup>7,8</sup>. <sup>1</sup>LESIA, Paris Observatory, F-92190 Meudon, France (thomas.widemann@obspm.fr); <sup>2</sup>University of Versailles-Saint-Quentin, Versailles, France; <sup>3</sup>LATMOS, 11 Boulevard d'Alembert, F-78280 Guyancourt, France; <sup>4</sup>SwRI, Boulder, CO; <sup>5</sup>Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA, 91109; <sup>6</sup>German Aerospace Center (DLR), Rutherfordstrasse 2, 12489 Berlin, Germany; <sup>7</sup>Planetary Science Institute, 1700 East Fort Lowell, Tucson, AZ 85719; <sup>8</sup>Dept. of Astronomy, Mount Holyoke College, South Hadley, MA 01075.

**Introduction:** The Venus Emissivity Mapper (VEM) is the first flight instrument specially designed to map the surface of Venus using the narrow atmospheric windows around 1  $\mu\text{m}$  [1]. VEM is proposed for the European Space Agency's M5/EnVision proposal in combination with a high-resolution radar mapper (*see Abstract #1937*). Mapping of Venus with VEx/VIRTIS using the 1.02  $\mu\text{m}$  thermal emission band can be viewed as a proof-of-concept for an orbital remote sensing approach to surface composition and weathering studies for Venus [2-7].

Thermal brightness on Venus' night side is mainly modulated by the lower clouds, imaged at 2.3  $\mu\text{m}$  by the Akatsuki IR2 camera [8]. Thanks to the circular polar orbit geometry of M5/EnVision, VEM has the unique capability to (1) better constrain the microphysics of the lower cloud particles in three spectral bands at 1.195, 1.310 and 1.510  $\mu\text{m}$  at a spatial resolution of  $\sim 10$  km, and (2) investigate short-timescale cloud dynamics and thus local wind speeds by tracking cloud features in both polar regions.

**Cloud parameters:** Global cloud layers ( $\sim 45$  to 70 km) drive the energy balance of the atmosphere and hence climate at the Venus' surface [9]. While much progress has been made since the early suggestion that the Venus clouds are  $\text{H}_2\text{O}$ - $\text{H}_2\text{SO}_4$  liquid droplets [10], several cloud parameters are still poorly constrained, particularly in the lower cloud layer and optically thicker polar regions [11-13]. Observations at small horizontal scales are of great importance to microphysical models of cloud and haze systems [14]. VEM has the capability to better constrain the microphysics (vertical, horizontal, time dependence of particle size distribution, or/and composition) of the lower cloud particles in three spectral bands at 1.195, 1.310 and 1.510  $\mu\text{m}$  at a spatial resolution of  $\sim 10$  km.

**Wind measurements:** Venus displays the best-known case of polar vortices evolving in a fast-rotating atmosphere. Few wind measurements exist in the polar region due to unfavorable viewing geometry of currently available observations. Cloud-tracking data indicate circumpolar circulation close to solid-body rotation. E-W winds decrease to zero velocity close to the poles. N-S circulation is marginal, with extremely variable morphology and complex vorticity patterns [15-17] (Fig. 1).

Circular polar orbit geometry would provide an unprecedented study of both polar regions within the same mission. VEM's pushbroom method will allow short-timescale cloud dynamics to be assessed, as well as local wind speeds, using repeated imagery at 90 minute intervals.

Tracking lower cloud motions as proxies for wind measurements at high spatial resolutions will greatly benefit modeling of the vortices' physics. Convective modeling demonstrates that there will be cloud-level




Figure 1 - Polar projection of South vortex morphology and cloud tracking zonal wind velocities near 1.74  $\mu\text{m}$  [17].

convection at high latitudes, so repeated imagery at 90 minute intervals will help constrain the time evolution of cloud-level convection as well as wave-generating dynamical instabilities [17]. This will also allow a direct comparison of the N-S wind regimes and their temporal evolution at several time scales.

**References:** [1] Helbert, J. et al. (2017) *SPIE Proc. Vol.10403, Infrared Remote Sensing and Instr. XXV*, San Diego, CA. [2] Mueller, N.T. et al. (2012), *Icarus* 217(2), 474-483. [3] D'Incecco, P. et al. (2016), *Pl. Space Sci.* 136, 25-33. [4] Helbert, J. et al. (2008), *Geophys. Res. Lett.* 35, L11201. [5] Gilmore, M.S., N. Mueller, and J. Helbert (2015) *Icarus* 254, 350-361. [6] Smrekar, S. et al. (2010) *Science* 328, 605-8. [7] Kappel, D. et al. (2016), *Icarus* 265, 42-62. [8] Gibney, E. (2016) *Nature* 532, 157-158. [9] Haus, R. et al., (2016) *Icarus*, 272, 178-205. [10] Young, A.T. (1973) *Icarus*, 18, 564-582. [11] Carlson, R.W et al. (1993), *Planet. Space Sci.*, 41, 411-485. [12] Wilson, C.F. et al. (2008), *J. Geophys. Res.*, 113, E00B13. [13] Barstow, J.K. et al. (2012), *Icarus* 217, 542-560. [14] McGouldrick, K. and Toon, O.B. (2007), *Icarus* 191, 1-24. [15] Sanchez-Lavega, A. et al. (2008), *Geophys. Res. Lett.* 35, L13204. [16] Luz, D. et al. (2011), *Science* 332, 577-580. [17] Garate-Lopez, I. et al. (2013) *Nature Geosc.* 6, 254-256.