



Modeling of aerosols from TIRVIM solar occultations onboard ExoMars/TGO

Mikhail Luginin, Nikolay Ignatiev, Anna Fedorova, Alexey Grigoriev, Alexey Shakun, Aleksander Trokhimovsky, Franck Montmessin, Oleg Korablev

► To cite this version:

Mikhail Luginin, Nikolay Ignatiev, Anna Fedorova, Alexey Grigoriev, Alexey Shakun, et al.. Modeling of aerosols from TIRVIM solar occultations onboard ExoMars/TGO. European Planetary Science Congress 2018, Sep 2018, Berlin, Germany. pp.EPSC2018-647. insu-04455551

HAL Id: insu-04455551

<https://insu.hal.science/insu-04455551>

Submitted on 13 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modeling of aerosols from TIRVIM solar occultations onboard ExoMars/TGO

Mikhail Luginin (1), N. Ignatiev (1), A. Fedorova (1), A. Grigoriev (1), A. Shakun (1), A. Trokhimovsky (1), F. Montmessin (2), and O. Koralev (1)

(1) Space Research Institute (IKI), Moscow, Russia (2) LATMOS, Guyancourt, France
(mikhail.luginin@phystech.edu)

The ExoMars Trace Gas Orbiter (TGO) is a joint ESA-Roscosmos mission to Mars that has been launched in March 2016. The aerobreaking phase has ended in February 2018 followed by the start of nominal scientific work on the near-circular 400 km orbit in April 2018. The Atmospheric Chemistry Suite (ACS) is a set of three spectrometers (NIR, MIR, and TIRVIM), capable to observe Mars atmosphere in solar occultations, nadir and limb geometry [1]. TIRVIM instrument is a Fourier-spectrometer operating in the 1.7 to 17 μm spectral range in solar occultation and nadir operation modes.

The main fraction of aerosols on Mars consists of mineral dust, while H_2O ice and CO_2 ice crystals are also encountered depending on the season and location. TIRVIM with its wide spectral range permits a spectral separation between dust and H_2O ice clouds particles. This work is dedicated to aerosols modeling in the solar occultation mode. Preliminary analysis of the transmission spectra retrieved from the TIRVIM solar occultation data shows presence of both dust and H_2O ice aerosol particles on some altitudes.

Acknowledgements

ExoMars is the space mission of ESA and Roscosmos. The ACS experiment is led by IKI Space Research Institute in Moscow. The project acknowledges funding by Roscosmos and CNES. Science operations of ACS are funded by Roscosmos and ESA. M. Luginin acknowledges the support from the Ministry of Education and Science of the Russian Federation, grant №14.W03.31.0017.

References

- [1] Koralev, O. et al. The Atmospheric Chemistry Suite (ACS) of Three Spectrometers for the ExoMars 2016 Trace Gas Orbiter. *Space Science Reviews*, 214(1), 7, 2018. <http://doi.org/10.1007/s11214-017-0437-6>