



HAL
open science

Analysis of Organic Molecules with Pyrolysis Gas Chromatography Mass Spectrometry in the Presence of Chlorides: Possible Implications for Measurements Performed with the SAM Experiment in Gale Crater

Cyril Szopa, Caroline Freissinet, A. Buch, Maëva Millan, B. Sutter, Michel Cabane, P. Glavin D., R. Navarro-Gonzalez, P. M. Mahaffy

► To cite this version:

Cyril Szopa, Caroline Freissinet, A. Buch, Maëva Millan, B. Sutter, et al.. Analysis of Organic Molecules with Pyrolysis Gas Chromatography Mass Spectrometry in the Presence of Chlorides: Possible Implications for Measurements Performed with the SAM Experiment in Gale Crater. 51st Lunar and Planetary Science Conference, Mar 2020, The Woodlands, United States. pp.LPI Contribution No. 2326, 2020, id.2219. insu-04460619

HAL Id: insu-04460619

<https://insu.hal.science/insu-04460619>

Submitted on 15 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ANALYSIS OF ORGANIC MOLECULES WITH PYROLYSIS GAS CHROMATOGRAPHY MASS SPECTROMETRY IN THE PRESENCE OF CHLORIDES : POSSIBLE IMPLICATIONS FOR MEASUREMENTS PERFORMED WITH THE SAM EXPERIMENT IN GALE CRATER.

C. Szopa^{1,2}, C. Freissinet¹, A. Buch³, M. Millan^{4,5}, B. Sutter⁶, M. Cabane¹, D.P. Glavin⁴, R. Navarro-Gonzalez⁷, and P. Mahaffy⁴, ¹LATMOS UVSQ/SU/CNRS, Guyancourt, France, ²Institut Universitaire de France, Paris, ³LGPM CentraleSupélec, Gif sur Yvette, France, ⁴NASA GSFC, Greenbelt, MD, ⁵Georgetown University, Washington D.C., ⁶Jacobs Technology, Inc., Johnson Space Center, National Aeronautics and Space Administration, Houston, Texas, ⁷Instituto de Ciencias Nucleares, UNAM, Mexico city, Mexico.

Introduction: Results from Checama, SAM, and APXS instruments onboard the Curiosity rover indicate that chlorides are present on Mars at Gale Crater, [1,2]. Even if no evaporitic layer of chlorides was found, these chemical species could represent a non-negligible fraction of chlorine bearing inorganic molecules present at the near surface of the crater floor. Laboratory analog studies have demonstrated that oxychlorine phases in Gale Crater material could serve as an O₂ and HCl source that causes the oxidation and chlorination of organic molecules present in samples analyzed by pyrolysis. SAM analysis detected chlorine bearing organic molecules in the Cumberland sample pyrolyzed at low temperatures (<400°C) [3,4]. This demonstrated that salts can interact with organic molecules present in a sample analyzed by pyrolysis. Total Cl from APXS analysis is always greater than SAM detected total Cl from oxychlorine phases indicating that chlorides likely dominate Cl in the Gale Crater materials. The goal of this work was to evaluate Mars analog chloride reactions with organic molecules to determine if the SAM detected chlorinated organics would form.

Moreover, other studies showed that other salts, as sulfates, can play a role of protection against the effect of oxychlorines during the pyrolysis [5,6,4]. Indeed, those salts start decomposing at higher temperatures than oxychlorines. Thus, organic molecules embedded in the salt matrix can be isolated from the gases released at low temperatures, preventing their reaction and alteration. Then, their protective role toward the effect of oxychlorine can be a key for detecting organic molecules indigenous to Mars.

The understanding of those reaction could also be a key to understand the organic molecule detected by SAM.

For these reasons, our team performed a systematic study in the laboratory of the possible influence of different chlorides on the analysis and detection of organic molecules. Then we performed pyrolysis gas chromatography mass spectrometry analysis, in order to assess their possible role in the analyses performed with the SAM experiment in Gale crater, and potentially future in situ experiments that would use such an analytical technique.

Experimental procedure: NaCl and MgCl₂ were used as the reference chlorides. Their thermal behavior was studied using thermal gravimetry (TG) and evolved gas analysis (EGA). More particularly their thermal decomposition was characterized as a function of the temperature up to about 900°C. Both chloride was then mixed with different organic molecules of interest for study of the Mars surface, i.e. polyaromatic hydrocarbons, amino acids, and carboxylic acids. Each mixture was submitted to two different pyrolytic treatments : i. a SAM like pyrolysis heating the sample from the ambient temperature up to 850°C at a 35°C.min⁻¹ rate; ii. a flash pyrolysis at 850°C. The products of pyrolysis were analyzed using gas chromatography mass spectrometry (GCMS) and compared.

Implications: The products of pyrolysis identified are compared to those detected with the SAM experiment. In particular, an emphasis will be done on the chlorinated organic molecules identified with SAM when analyzing the Cumberland sample collected in the Yellow Knife bay area early in the mission. Conclusions about the possible influence of chlorides on GCMS analyses of Mars surface samples will be done.

Acknowledgments: NASA Mars Exploration Program supported for the development and operations of SAM. The French National Space Agency, the Centre National d'Études Spatiales (CNES), supported the development and operations of the SAM gas chromatograph.

References: [1] Sutter et al., , JGR, , 1 (2017). [2] N.H. Thomas et al., GRL, 46 (2019) 10754-10763. [3] C. Freissinet et al., JGR 120 (2015) 2014JE004737. [4] C. Szopa et al., Astrobiology, (In press). ; [5] P. Francois et al., JGR., 121 (2016) 61-74.[6] M.T.J. Lewis et al., Astrobiology, (2015).