



HAL
open science

Main ions in the thermosphere of exoplanets, focussing on the transition between super-Earths and mini-Neptunes

Jérémy Bourgalais, Nathalie Carrasco

► **To cite this version:**

Jérémy Bourgalais, Nathalie Carrasco. Main ions in the thermosphere of exoplanets, focussing on the transition between super-Earths and mini-Neptunes. EPSC-DPS Joint Meeting 2019, Sep 2019, Geneva, Switzerland. pp.EPSC-DPS2019-1364-1. insu-04462240

HAL Id: insu-04462240

<https://insu.hal.science/insu-04462240>

Submitted on 16 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Main ions in the thermosphere of exoplanets, focussing on the transition between super-Earths and mini-Neptunes

Jérémy Bourgalais (1) and Nathalie Carrasco (1)

(1) LATMOS/IPSL, UVSQ, Université Paris-Saclay, UPMC Université Paris 06, CNRS, Guyancourt, France
(Jeremy.bourgalais@latmos.ipsl.fr)

Abstract

Therospheres of exoplanets are submitted to the harsh UV and X-ray radiations of their host star. In the present work we focus on ion chemistry occurring in super-Earths and mini-Neptunes. We simulate in the laboratory the effect of Far-UV on gas-mixtures varying from H₂-rich to H₂-poor conditions and we analyze the ions by *in situ* mass spectrometry.

1. Introduction

Therospheres of exoplanets are photochemically active layers, characterized by an efficient photolysis of the main components of the atmospheres. Vertical transport in therospheres is moreover governed by molecular diffusion, also contributing to strongly modify the composition profiles of the atmospheres. On Mars, with an atmosphere based on CO₂ and N₂, the efficient photolysis of CO₂ leads to an increase of CO mixing ratio, up to several tens percent's compared to CO₂ [1]. The impact on the thermal profile of the atmosphere is determining. On Mars the progressive enrichment in CO compared to CO₂ according to altitude limits the heating of the therosphere, as CO is a less efficient UV absorber than CO₂ [2]. Predicting and understanding the temperature and density profiles of therospheres of exoplanets will therefore constrain atmospheric escape and photoevaporation processes, and a fortiori the evolution of atmospheres of exoplanets.

Photolysis in therospheres involves harsh UV and X-ray stellar radiations and produces complex chemical environments coupling neutral and ion chemistry. Ion-chemistry has been shown to be largely underestimated and poorly known when the Cassini-Huygens space mission unveiled the complex ion and neutral composition of the therosphere of

the largest satellite of Saturn, Titan [3]. Its atmosphere is based on N₂ and CH₄, and nitrogen chemistry was found to be particularly important in Titan's therosphere. We learned with the study of this body that ion-neutral chemistry was rich and efficient and could lead to unexpected abundant chemical species.

With the data of the California-Kepler Survey two populations of super-Earths and mini-Neptunes could be distinguished [4]. Super-Earths, with radii lower than 1.5 R_⊕ are rocky planets with thin atmospheres mainly made of heavy molecules like N₂ and CO₂. Mini-Neptunes with masses between 2 and 3 R_⊕ have on the opposite extended H₂-based atmospheres.

The aim of the present work is to characterize the main positive ions that can be expected in the therospheres of super-Earths and mini-Neptunes. For this purpose we simulated experimentally the ion-neutral chemistry initiated by Far-UV radiations on a gas mixtures mimicking the main possible composition of therospheres of super-Earths and mini-Neptune. The ion species are analyzed *in situ* by mass spectrometry.

2. Methods

Far-UV photons mimicking the stellar UV field were generated by using a gas-discharge type Far-UV source coupled windowless to our photochemical reactor [5]. We irradiated at 73.6 nm (15.6 eV) by using the resonant emission lines of Ne I and Ar I, respectively, with a photon flux of ca. 1.5×10^{14} ph.s⁻¹.cm⁻².

Gas mixtures were chosen to be representative of H₂-poor atmospheres and H₂-rich atmospheres. As the photolysis of CO₂ mainly produces CO, and that CO

is more efficient to drive further photochemistry, we have chosen CO as a proxy of both CO₂ and CO in our experiments. In order to build a consistent experimental series, we have chosen to only modify the H₂ content in our experimental conditions. N₂ mixing ratio was therefore arbitrary chosen as equal to CO mixing ratio, and H₂ mixing ratios increased from 1% up to 96%.

The gas composition was monitored in steady state by using a high sensitivity (<1ppm) quadrupole mass spectrometer (Hiden Analytical EQP 200 QMS) allowing *in situ* measurements of the positive ions.

3. Results

H₂ mixing ratios were chosen at 1%, 33% and 96%. The main ions detected are shown on Figure 1. The first striking result is that in all conditions the main ions are the same: H₃O⁺ at m/z 19 and HCO⁺ and N₂H⁺ at m/z 29, even in a H₂-based neutral gas mixture. In the H₂-based condition (H₂ mixing ratio of 96%), H₃⁺ at m/z 3 appears as an additional important ion contributor.




Figure 1: Ions produced after irradiation at 73 nm of H₂/CO/N₂ gas mixtures. H₂ mixing ratios was chosen at 1%, 33% and 96%.

Primary ions produced by photoionization at 73nm are respectively H₂⁺ at m/z 2, N₂⁺ and CO⁺ at m/z 28. Those primary ions react with H₂, through ion-molecule efficient reactions, leading to the stable protonated ions observed. The detection of the ion H₃O⁺ is a sensitive probe of the presence of H₂O in the reactive medium: as a photoproduct and/or as contaminant.

4. Summary and Conclusions

We highlight that the main ions observed in our H₂ rich (96%) and H₂ poor (1%) conditions are similar and corresponds to HCO⁺ and N₂H⁺. We used CO as a proxy for CO and CO₂. We can extrapolate that HCO₂⁺ would similarly be observed in the presence of CO₂. Despite the drastic differences in neutral compositions of the atmospheres of super-Earths and mini-Neptunes, our work suggests that rather similar ion contributions could be expected in their thermospheres.

Acknowledgements

This research was supported by the ERC Starting Grant PRIMCHEM, grant agreement n°636829.

References

- [1] Moudén Y. and Connell J.C. *Icarus*, 188, 18, 2007.
- [2] Gonzalez-Galindo, F., Forget, F., Angelats, M., Coll, I. and Lopez-Valverde, M. A. *Lecture Notes and Essays in Astrophysics*, 3, 151, 2008.
- [3] Waite, J. H., Young, D. T., Cravens, T. E., et al. *Science*, 316, 870, 2007.
- [4] Fulton F., Petigura E., Howard A., et al. *Astronomical Journal*, 154(3), 109, 2017.
- [5] Tigrine S., Carrasco N., Vettier L., Cernogora G. *Journal of Physics D: Applied Physics*, 49, 395202, 2016