



HAL
open science

Atmospheric noble gas isotope and bulk K/U ratios as a constraint on the early evolution of Venus and Earth

H. Lammer, M. Scherff, M Leitzinger, P. Odert, D. Kubyshkina, C. Burger, C. Johnstone, T. Maindl, M. Güdel, N. Tosie, et al.

► To cite this version:

H. Lammer, M. Scherff, M Leitzinger, P. Odert, D. Kubyshkina, et al.. Atmospheric noble gas isotope and bulk K/U ratios as a constraint on the early evolution of Venus and Earth. EPSC-DPS Joint Meeting 2019, Sep 2019, Geneva, Switzerland. pp.EPSC-DPS2019-1550-1. insu-04469246

HAL Id: insu-04469246

<https://insu.hal.science/insu-04469246v1>

Submitted on 20 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Atmospheric noble gas isotope and bulk K/U ratios as a constraint on the early evolution of Venus and Earth

H. Lammer (1), **M. Scherf** (1), M. Leitzinger (1,2), P. Odert (1,2), D. Kubyskhina (1), C. Burger (3), C.P. Johnstone (3), T. Maindl (3), M. Güdel (2), N. Tosi (4,5), E. Marcq (6), N.V. Erkaev (7,8), L. Fossati (1)

(1) Space Research Institute, Austrian Academy of Sciences, Graz, Austria, (2) Institute of Physics/IGAM, University of Graz, Austria, (3) Institute of Astrophysics, University of Vienna, Vienna, Austria, (4) Institute of Planetary Research, Department of Planetary Physics, DLR, Berlin, Germany, (5) Department of Astronomy and Astrophysics, Berlin Institute of Technology, Germany, (6) LATMOS, Université de Versailles Saint-Quentin-en-Yvelines, Gyancourt, France, (7) Institute of Computational Modelling SB RAS, Krasnoyarsk, Russian Federation, (8) Siberian Federal University, Krasnoyarsk, Russian Federation (helmut.lammer@oeaw.ac.at)

Abstract

While Venus and Earth were accumulating mass within the solar nebular these protoplanets also captured significant hydrogen dominated atmospheres by picking up gas from the circumstellar disk during the formation of the Solar System (e.g. [1]). These primordial atmospheres were then quickly lost by hydrodynamic escape after the disk dissipated. After a short but efficient boil-off phase the EUV-driven hydrodynamic flow of H-atoms dragged heavier elements with it at different rates, leading to changes in their isotopic and elemental ratios [2,3,4], which is reflected in the present-day atmospheric noble gas isotope and elemental ratios of Venus and Earth. Depending on the disk lifetime and the initial composition $^{36}\text{Ar}/^{38}\text{Ar}$, $^{20}\text{Ne}/^{22}\text{Ne}$ and bulk K/U ratios observed for both planets can be best explained if the Sun was born between a weakly and moderately active star and if Venus and Earth had grown to ~85-100% and ~53-58%, respectively, of their current masses by the time the nebula gas dissipated approximately 3.5 Myr after formation of the Sun. If proto-Earth accreted its mass from up to 40% carbonaceous chondritic-like material [5] then the planet must have been grown to about 80% of its final mass as long as it was surrounded by the escaping primordial atmosphere (~7 Myr). Our results are therefore in agreement with a fast accretion of thermally processed disk material planetary embryos, with Hafnium-Tungsten chronometric fast accretion scenarios of the proto-Earth (e.g. [6]), and a noble gas origin based on a mixture of primitive meteoroids and a small remnant of the proto-solar nebula [7].

References

- [1] Stökl, A., Dorfi, E.A., Johnstone, C.P., Lammer, H.: Dynamical Accretion of Primordial Atmospheres around Planets with Masses between 0.1 and 5 M in the Habitable Zone, *Astrophys. J.*, 825,86, 2016.
- [2] Hunten, D.M., Pepin, R.O., Walker, J.C.G.: Mass fractionation in hydrodynamic escape, *Icarus*, 69, 532-49, 1987.
- [3] Zahnle, K., Pollack, J.B., Kasting, J.F.: Mass fractionation of noble gases in diffusion-limited hydrodynamic hydrogen escape, *Icarus*, 84, 502–27, 1990.
- [4] Odert, P, Lammer, H, Erkaev, N.V, Nikolaou, A., Lichtenegger, H.I.M., Johnstone, C.P., Kislyakova, K.G., Leitzinger, M., Tosi, N.: Escape and fractionation of volatiles and noble gases from Mars-sized planetary embryos and growing protoplanets, *Icarus*, 307, 327–46, 2018.
- [5] Schiller, M., Bizzarro, M., Fernandes, V.A.: Isotopic evolution of the protoplanetary disk and the building blocks of Earth and the Moon, *Nature*, 555, 507–10, 2018.
- [6] Yu, G., Jacobsen, S.B.: Fast accretion of the Earth with a late Moon-forming giant impact, *Proceedings of the National Academy of Science*, 108, 17604–9, 2011.
- [7] Marty, B.: The origins and concentrations of water, carbon, nitrogen and noble gases on Earth. *Earth and Planetary Science Letters*, 313, 56–66, 2012.