



Using satellite probabilistic estimates to assess modeled relative humidity: application to a NWP model

Chloé Radice, Hélène Brogniez, Pierre-Emmanuel Kirstetter, Philippe Chambon

► To cite this version:

Chloé Radice, Hélène Brogniez, Pierre-Emmanuel Kirstetter, Philippe Chambon. Using satellite probabilistic estimates to assess modeled relative humidity: application to a NWP model. EGU23, the 25th EGU General Assembly, Apr 2023, Vienna, Austria. pp.id. EGU-13778, 10.5194/egusphere-egu23-13778 . insu-04475800

HAL Id: insu-04475800

<https://insu.hal.science/insu-04475800>

Submitted on 5 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Using satellite probabilistic estimates to assess modeled relative humidity : application to a NWP model

C. Radice (chloe.radice@latmos.ipsl.fr), H. Brogniez, P-E. Kistetter, P. Chambon

Deterministic method : is the simulated value within a fixed interval around the mean observed value ?



Probabilistic method : uses the full estimated distribution to determine the probability of the simulated value being an over- or underestimation



Probabilistic RH estimates from microwave sounder **SAPHIR** aboard satellite Megha-Tropiques

Simulated RH by the Numerical Weather Prediction model **ARPEGE** developed by Météo France

Both methods were applied to 3 months worth of the ARPEGE-SAPHIR colocated dataset (April-May-June 2018)



Deterministic assessment: slight humid bias over the ITCZ

Probabilistic assessment: overall overestimation with local patches of underestimation

Even when the simulated values are close to the mean estimation, they actually correspond to the extreme ends of the entire subgrid distribution.

→ Complementary to a deterministic approach + adds contrast within the deterministic interval



Using satellite probabilistic estimates to assess modeled relative humidity : application to a NWP model

C. RADICE (CHLOE.RADICE@LATMOS.IPSL.FR), H. BROGNIEZ, P-E. KIRSTETTER, P. CHAMBON



Context


Datasets

ARPEGE at Météo-France

- Numerical weather forecasting model operated by Météo-France
- Forecasts initialized with a four dimensional variational system with 6h windows (00:00, 06:00, 12:00 and 18:00 UTC)
- For this study : 6-hourly instantaneous simulated RH fields at 6 hour lead time for the months April, May and June 2018.
- The resolution of the model grid used is $0.25^\circ \times 0.25^\circ$

SAPHIR microwave sounder

- 6 channels sampling the 183 GHz water vapour absorption line
- Onboard Indo-French (ISRO x CNES) satellite Megha-Tropiques
- Scans the tropical belt (between 30°N and 30°S)
- Retrieval scheme for a profile of RH over 6 layers (100 to 950 hPa)
- For each layer : expected **RH** as well as its **PDF** to account for uncertainty in the retrieval ([Sivira et al 2015](#), [Brogniez et al 2016](#)).





Methodology


Assessing forecasts with satellite observations

Colocation

- ± 30 min time frame around the ARPEGE simulation
- Atmospheric levels of the simulations are aggregated together to mimic the vertical resolution of the SAPHIR RH profiles (6 pressure layers)

Our probabilistic method

- Convoluting the PDF of every footprints within each gridbox of ARPEGE
- Projecting the simulated value (SV) onto the distribution



Deterministic comparison


- For every gridbox of ARPEGE, spatial average of SAPHIR $\overline{\text{RH}}$
- Confronting the simulated value (SV) to this averaged value



Methodology Application


Deterministic comparison

- \bar{RH} computed from the PDF
 - + Confidence interval arbitrarily set to $\pm 15\%RH$
(defined from a global evaluation of the retrieval using radiosoundings)
- Difference : $D = SV - \bar{RH}$
- Example : ARPEGE $SV = 30\%RH \rightarrow D = 11\%RH$



Our probabilistic method

- PDF → Cumulative Density Function (CDF)
- $CDF(SV) = \mathcal{P}(RH \leq SV) = P$
- Situates SV in the estimated CDF within or outside the inter-quartile 0.25-0.75
- Example : $P=0.84$
 - probability of 84% that the true RH estimate is lower than SV
 - SV falls within the upper quarter of the distribution ($P > 0.75$)
 - ⇒ High probability that the model overestimates the RH




Motivations

Comparing to the same deterministic interval does not account for the shape of the whole distribution.

IQR = Interquartile Range ($Q_3 - Q_1$ or $X_{P(X)=0.75} - X_{P(X)=0.25}$) of the subgrid distributions

$IQR < 15\%RH \rightarrow$ the subgrid distribution has a smaller uncertainty than the $\pm 15\%RH$ deterministic interval
 \rightarrow the $\pm 15\%RH$ error works

$IQR > 15\%RH \rightarrow$ the subgrid distribution is too wide to fit the $\pm 15\%RH$ deterministic interval
 \rightarrow need to have a wider error than $\pm 15\%RH$







Results

Maps

One time step, one atmospheric layer



01/04/2018 – 00UTC – 400 to 600 hPa



- Same structures of biases highlighted by the two methods (see zoomed areas)
- Overall, with the deterministic method, ARPEGE SV close to the mean estimate (within the $\pm 15\%$ RH interval)
- The probabilistic method shows very contrasting patterns (<0.25 or >0.75)

3-month period, one layer

April to June 2018 – 400 to 600 hPa



- The deterministic method only highlights a slight moist bias (\bar{D} between 15 and 30 %RH) in the Intertropical Convergence Zone (ITCZ)
- The probabilistic method shows a very high probability (>0.8) that the model overestimates the atmospheric RH for this pressure layer
- Some patches of probable underestimation of RH (<0.2) appear only with this method (above the south of Africa, the Indian Ocean, equatorial Pacific Ocean, etc.)




Results


Distributions for the whole tropical belt (30°N-30°S)

1 time step, 1 atmospheric layer


01/04/2018 – 00UTC – 400 to 600 hPa



Distribution of the deviations from mean (D)
ARPEGE Spring 2018



Distribution of the associated percentiles (P)
ARPEGE Spring 2018



- Grey area not symmetrical around the $y=x$ line
→ [0.25;0.75] interval not centred around the mean
- A fixed confidence interval (ex. $\pm 15\text{ %RH}$) is not suitable and the knowledge of the spread of the retrieval (PDF/CDF) better quantifies the assessment of the model

- When SV outside $\pm 15\text{ %RH}$, P in majority in the extreme 5% of the distributions (<0.5 or >0.95)
- Added information content in the probabilistic results (wide distributions) wrt the deterministic method (narrower distributions with few extreme values)





Sharing is encouraged




Conclusion

- The probabilistic method adds significant information to the comparisons, adapts to each situation and allows a both robust and precise diagnosis.
- This probabilistic method shows promising results on this data set but is perfectly adaptable to any other variable, model and reference PDF
- Check out [radice et al., 2022](#) for more details !



Scan me to get to
Radice et al., 2022



Atmospheric water cycle studying
satellite Megha-Tropiques
© CNES/ISRO

Grid of the numerical weather forecasting model ARPEGE
© Météo-France

Chloé Radice (chloe.radice@latmos.ipsl.fr) – LATMOS/IPSL – Université Paris-Saclay