



HAL
open science

Detecting Venus' volcanic gas plumes with VenSpec-H

Séverine Robert, Caterina Macovenco, Maxence Lefèvre, Colin Wilson,
Emmanuel Marcq, Bruno Bézard, Jörn Helbert, Ann Carine Vandaele

► **To cite this version:**

Séverine Robert, Caterina Macovenco, Maxence Lefèvre, Colin Wilson, Emmanuel Marcq, et al..
Detecting Venus' volcanic gas plumes with VenSpec-H. European Planetary Science Congress 2021,
Sep 2021, Online, United States. pp.EPSC2021-678, 10.5194/epsc2021-678 . insu-04476661

HAL Id: insu-04476661

<https://insu.hal.science/insu-04476661>

Submitted on 25 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Europlanet Science Congress 2021

Virtual meeting

13 September – 24 September 2021

[Home](#) > EPSC2021-678

EPSC Abstracts

Vol. 15, EPSC2021-678, 2021

<https://doi.org/10.5194/epsc2021-678>

European Planetary Science Congress 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Detecting Venus' volcanic gas plumes with VenSpec-H

Séverine Robert et al.

The VenSpec-H instrument is part of the EnVision payload which is currently being evaluated by ESA for mission selection. EnVision is a medium class mission to determine the nature and current state of geological activity on Venus, and its relationship with the atmosphere, to understand how Venus and Earth could have evolved so differently.

VenSpec-H is part of the VenSpec suite [1], including also an IR mapper and a UV spectrometer [2] suite. The science objectives of this suite are to search for temporal variations in surface temperatures and tropospheric concentrations of volcanically emitted gases, indicative of volcanic eruptions; and study surface-atmosphere interactions and weathering by mapping surface emissivity and tropospheric gas abundances. Recent and perhaps ongoing volcanic activity has been inferred in data from both Venus Express and Magellan. Maintenance of the clouds requires a constant input of H₂O and SO₂. A large eruption would locally alter the composition by increasing abundances of H₂O, SO₂ and CO and perhaps decreasing D/H ratio. Observations of changes in lower atmospheric SO₂, CO and H₂O vapour levels, cloud level H₂SO₄ droplet concentration, and mesospheric SO₂, are therefore required to link specific volcanic events with past and ongoing observations of the variable and dynamic mesosphere, to understand both the importance of volatiles in volcanic activity on Venus and their effect on cloud maintenance and dynamics.

To contribute to this investigation, VenSpec-H is designed to measure H₂O and HDO contents in the first scale height of Venus' atmosphere and probe H₂O, HDO, CO, OCS, SO₂ in the 30 to 40 km altitude range [3-8]. To assess the performances of our instrument at detecting volcanic eruptions, we defined a couple of scenarios of plume releases and simulated the corresponding spectra.

To simulate the vertical transport of the plume and the horizontal advection by the dynamics, the LMD Venus Mesoscale Model [9] is used, based on the WRF dynamical core and the Venus IPSL radiative transfer. The domain is focused on Imdr Regio, where VIRTIS observed a hotspot anomaly possibly linked to volcanic activity [10]. Tracers were added to the model representing H₂O, CO and SO₂. The chemistry and photodissociation sources and sinks are modelled by a linear relaxation of the tracer abundance toward a value representative of the deep atmosphere with a characteristic time. The deep atmosphere abundance is set to 130 ppm for SO₂, 30 ppm for H₂O and 20 ppm for CO. The relaxation time is set to 100 years for SO₂, to 1000 years for CO and to 1 week, 1 month and 1 year for H₂O representing the uncertainty of the chemistry timescale in that region. Several configurations are considered for the plume, an idealised set-up where the elevation height is fixed and the outgassing abundance is constant inside the plume and in time, and a more realistic set-up where a temperature and outgassing anomalies at the surface are prescribed is ongoing testing.

The radiances of the nightside atmosphere of Venus originate from the thermal emission of the surface and atmosphere. The spectra were simulated from 1 to 2.5 microns, using ASIMUT-ALVL, a line-by-line radiative transfer code developed at BIRA-IASB [11]. CO₂, H₂O, HDO, CO, SO₂, and OCS, as well as aerosols were included.

The performances of the instrument will be described in terms of its capabilities to detect small variations in the atmosphere.

References

- [1] J. Helbert et al., "The VenSpec suite on the ESA EnVision mission to Venus", *Proc. SPIE 11128, Infrared Remote Sensing and Instrumentation XXVII*, 1112804 (2019).
- [2] E. Marcq et al., "Instrumental requirements for the study of Venus' cloud top using the UV imaging spectrometer VeSUV", *Advances in Space Research*, 68 (2021) 275-291.
- [3] B. Bézard et al., "Water vapor abundance near the surface of Venus from Venus Express/VIRTIS observations," *Journal of Geophysical Research, Planets*, 114 (2009) E00B39.
- [4] B. Bézard et al., "The 1.10- and 1.18- μ m nightside windows of Venus observed by SPICAV-IR aboard Venus Express," *Icarus*, 216 (2011) 173-183.
- [5] A. Fedorova et al., "The CO₂ continuum absorption in the 1.10-and 1.18- μ m windows on Venus from Maxwell Montes transits by SPICAV IR onboard Venus Express," *Planetary and Space Science*, 113 (2015) 66-77.

[6] E. Marcq et al., "Remote sensing of Venus' lower atmosphere from ground-based IR spectroscopy: latitudinal and vertical distribution of minor species," *Planetary and Space Science*, 54 (2006) 1360-1370.

[7] E. Marcq et al., "A latitudinal survey of CO, OCS, H₂O, and SO₂ in the lower atmosphere of Venus: spectroscopic studies using VIRTIS-H," *Journal of Geophysical Research, Planets*, 113 (2008) E00B07.

[8] E. Marcq et al., "Evidence for SO₂ latitudinal variations below the clouds of Venus", *Astronomy & Astrophysics*, 648 (2021) L8.

[9] M. Lefèvre, A. Spiga and S. Lebonnois, "Mesoscale modeling of Venus' bow-shape waves", *Icarus*, 335 (2019) 113376.

[10] E. Smrekar, et al., "Recent Hotspot Volcanism on Venus from VIRTIS Emissivity", *Data. Science*, 328 (2010) 605.

[11] A.C. Vandaele, M. Kruglanski and M. De Mazière, "Modeling and retrieval of atmospheric spectra using ASIMUT", *Proc. of the First 'Atmospheric Science Conference'*, ESRIN, Frascati, Italy, 2006.

How to cite: Robert, S., Macovenco, C., Lefèvre, M., Wilson, C., Marcq, E., Bézard, B., Helbert, J., and Vandaele, A. C.: Detecting Venus' volcanic gas plumes with VenSpec-H, European Planetary Science Congress 2021, online, 13–24 Sep 2021, EPSC2021-678, <https://doi.org/10.5194/epsc2021-678>, 2021.



Contact

[Imprint](#)

[Contact](#)



[Data protection](#)

© 2021 Copernicus GmbH

Header image © Europlanet Society