



HAL
open science

Three-dimensional modeling of the O₂(1Δ) dayglow and implications for ozone in the middle atmosphere.

Mouhamadou Diouf, Franck Lefèvre, Alain Hauchecorne, Jean-Loup Bertaux

► **To cite this version:**

Mouhamadou Diouf, Franck Lefèvre, Alain Hauchecorne, Jean-Loup Bertaux. Three-dimensional modeling of the O₂(1Δ) dayglow and implications for ozone in the middle atmosphere.. EGU General Assembly 2024, Apr 2024, Vienna and Online, Austria. pp. EGU24-1897, 10.5194/egusphere-egu24-1897. insu-04501100

HAL Id: insu-04501100

<https://insu.hal.science/insu-04501100v1>

Submitted on 3 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU24-1897, updated on 03 Jun 2024

<https://doi.org/10.5194/egusphere-egu24-1897>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Three-dimensional modeling of the $O_2(^1\Delta)$ dayglow and implications for ozone in the middle atmosphere.

Mouhamadou Diouf, Franck Lefevre, Alain Hauchecorne, and Jean-Loup Bertaux

LATMOS, CNRS, Sorbonne Université, Université Paris-Saclay, Paris, France (mouhamadou.diouf@latmos.ipsl.fr)

Future space missions dedicated to measuring CO_2 on a global scale can make advantageous use of the O_2 band at $1.27 \mu m$ to retrieve the air column. The $1.27 \mu m$ band is close to the CO_2 absorption bands at 1.6 and $2.0 \mu m$, which allows a better transfer of the aerosol properties than with the usual O_2 band at $0.76 \mu m$. However, the $1.27 \mu m$ band is polluted by the spontaneous dayglow of the excited state $O_2(^1\Delta)$, which must be removed from the observed signal.

We investigate here our quantitative understanding of the $O_2(^1\Delta)$ dayglow with a chemistry-transport model. We show that the previously reported -13% deficit in $O_2(^1\Delta)$ dayglow calculated with the same model is essentially due to a -20 to -30% ozone deficit between 45 - 60 km. We find that this ozone deficit is due to excessively high temperatures ($+15$ K) of the meteorological analyses used to drive the model in the mesosphere.

The use of lower analyzed temperatures (ERA5), in better agreement with the observations, slows down the hydrogen-catalyzed and Chapman ozone loss cycles. This effect leads to an almost total elimination of the ozone and $O_2(^1\Delta)$ deficits in the lower mesosphere. Once integrated vertically to simulate a nadir measurement, the deficit in modeled $O_2(^1\Delta)$ brightness is reduced to $-4 \pm 3\%$. This illustrates the need for accurate mesospheric temperatures for a priori estimations of the $O_2(^1\Delta)$ brightness in algorithms using the $1.27 \mu m$ band.