

Scientific interest for a UV spectrometer to study Uranus and its satellites

Jean-Yves Chaufray, Laurent Lamy, François Leblanc, V Hue, Franck Montmessin, Eric Quémerais, Emmanuel Marcq, S Guerlet, G Milcareck, M Barthelemy

▶ To cite this version:

Jean-Yves Chaufray, Laurent Lamy, François Leblanc, V Hue, Franck Montmessin, et al.. Scientific interest for a UV spectrometer to study Uranus and its satellites. Uranus Flagship: Investigations and Instruments for Cross-Discipline Science Workshop, Jul 2023, Pasadena, United States. pp.LPI Contribution No. 2808, 2023, id. 8052. insu-04501879

HAL Id: insu-04501879 https://insu.hal.science/insu-04501879v1

Submitted on 12 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Scientific interest for a UV spectrometer to study Uranus and its satellites. J-Y. Chaufray¹, L. Lamy^{2,3}, F.Leblanc¹, V. Hue³, F. Montmessin¹, E.Quémerais¹, E. Marcq¹, S. Guerlet³, G. Milcareck¹, M. Barthelemy⁵ Affiliation (¹ LATMOS/IPSL, CNRS, Paris-Saclay, Sorbonne Université, Guyancourt, France, <u>chaufray@latmos.ipsl.fr</u>), ²LESIA, Observatoire de Paris, PSL, Meudon, France, ³ Aix-Marseille Université, CNRS, CNES, Institut Origines, LAM, Marseille, France, ⁴LMD, Paris, France, ⁵IPAG, Grenoble, France.

Introduction: Uranus and its satellites remain poorly known and were only explored during the flyby of Voyager-2 in 1986 and a few observations from Earth. New UV observations of Uranus and its satellites on a future orbiter could provide important information about: i) the interaction of the Uranian ionosphere and its magnetosphere, ii) its atmospheric structure and composition and iii) the icy surface of its satellites.

Aurorae, airglow and electroglow: The magnetosphere of Uranus is very different from the other giant planets magnetosphere due to its highly tilted magnetic field [1] and present large diurnal/seasonal variations. Like the other giant planets, the UV auroral emissions are confined near the magnetic poles [2] and dominated by the H (Lyman- α) and H₂ (Lyman and Werner bands).

The H Lyman- α and emissions from H₂ were first measured by the UV spectrometer of Voyager-2 [3], and more recently re-observed by HST/STIS [4]. The Lyman- α emission was found to be produced by the resonant scattering of the solar photons by atomic hydrogen in the corona, by resonant scattering of the interplanetary Lyman-a photons, by Rayleigh scattering of the solar photons by the atmospheric H_2 [3,5], and a possible additional unknown internal source. The Rayleigh scattering was not expected but suggested by the presence of a Raman emission near 128 nm and requires a depletion in hydrocarbons above 0.5 mbar possibly due to a very low eddy diffusion coefficient [3,6]. However, this low value has never been confirmed from later observations [7] and our estimate of the process using more recent atmospheric models of Uranus [8] suggests it could have been overestimated by [3]. The possible additional unknown internal source could be due to low energy impacts ("electroglow") [1].

Atmosphere of Uranus: Planetary atmosphere can be investigated remotely in the UV either by reflectance measurement from nadir observations or by stellar occultations. Occultations of γ -Pegasi and the sun were obtained by Voyager-2/UVS to measure the vertical profiles of the absorbing species [9] leading to larger eddy diffusion compared to [3]. On Uranus, the atmospheric UV extinction should be mostly due to H₂ (absorption and Rayleigh and Raman scattering), and hydrocarbons species below 200 nm (Fig. 1) and by H₂ Rayleigh and Raman scattering and aerosol (from stratospheric haze) between 200 and 400 nm [10].




Fig. 1 UV extinction cross sections from the major gas species in the atmosphere of Uranus. Based on https://phidrates.space.swri.edu, and [11] for Rayleigh and Raman extinction by H_2 (at 100 K).

Surfaces of satellites:

The icy surface of the satellites of Uranus should have an albedo similar to other icy satellites in the solar system with a low albedo below 170 nm and a large albedo above 170 nm. Observations done by HTS/FOS instrument find a rather flat albedo for Ariel, Titania and Oberon between 220 and 1000 nm but with a possible small absorption near 280 nm and decreasing with increasing distance to Uranus. This absorption could be due to entrapped OH gas but needs to be confirmed [12].

In our presentation we will discuss the measurements from a UV spectrometer that would help on these three subjects and present our Monte Carlo model of Rayleigh/Raman extinction to study some UV emission lines.

References:

[1] Broadfoot, et al., (1986), *Science*, 233, 74 [2] Lamy. et al. (2017), JGR, 122, 3997-4008.[3] Yelle et al; (1987), GRL, 14, 483-486, [4] Barthelemy et al. (2014), Icarus, 239 [5] Emerich et al. (1993), PSS, 41, 363, [6], Yelle et al. (1989), Icarus, 77, 439-456, [7] Encrenaz et al. (1998), A&A, 333, L43-L46, [8] Moses et al., (2020), Philos. Trans. R. Soc. A. [9] Bishop et al. (1990), Icarus, 88, 448-464, [10] Cochran et al. (1990), Icarus, 83, 93-101, [11] Ford and Browne (1973), At. Dat., 5, 305-313, [12] Hendrix et al. (2012), *in the science of solar system ices* (Vol. 356), Astrophysics and Space Science Library.

Acnkowledgements:

We acknowledge support from CNES and the CNRS/INSU programs of planetology (PNP- and heliophysics (PNST).