

Stratospheric impact of the anomalous 2023 Canadian wildfires

Sergey Khaykin, Sophie Godin-Beekmann, Slimane Bekki, Florent Tencé, Mehdi Meziane, Béatrice Josse, Sophie Pelletier, Qiaoyun Hu, Philippe Goloub, Alexandra Laeng

▶ To cite this version:

Sergey Khaykin, Sophie Godin-Beekmann, Slimane Bekki, Florent Tencé, Mehdi Meziane, et al.. Stratospheric impact of the anomalous 2023 Canadian wildfires. EGU General Assembly 2024, Apr 2024, Vienna, Austria. pp.EGU24-5686, 10.5194/egusphere-egu24-5686. insu-04507984

HAL Id: insu-04507984 https://insu.hal.science/insu-04507984v1

Submitted on 4 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



EGU24-5686, updated on 04 Jun 2024 https://doi.org/10.5194/egusphere-egu24-5686 EGU General Assembly 2024 © Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Stratospheric impact of the anomalous 2023 Canadian wildfires

Sergey Khaykin¹, Sophie Godin-Beekmann¹, Slimane Bekki¹, Florent Tence¹, Mehdi Meziane², Beatrice Josse², Sophie Pelletier², Qiaoyun Hu³, Philippe Goloub³, and Alexandra Laeng⁴ ¹LATMOS, CNRS/INSU, UVSQ, Sorbonne university, Guyancourt, France (sergey.khaykin@latmos.ipsl.fr) ²CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France ³LOA – Laboratoire d'Optique Atmosphérique, CNRS, University of Lille, Lille, France ⁴Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

The frequency of extreme wildfires has increased as a response to the regional and global warming trends and there is an emerging realization of their impact on climate through emissions of smoke aerosols into the stratosphere. The 2023 wildfire season in Canada was unprecedented in terms of its duration, burned area and cumulative fire power, rendering it the most destructive ever recorded.

Here we use various satellite observations (TROPOMI, OMPS-LP, OMPS-NM, MLS, CALIPSO, SAGE III) to quantify the stratospheric emissions of smoke aerosols and carbon monoxide by the 2023 Canadian wildfires and to characterize the long-range transport of smoke plumes in the stratosphere. Using multiwavelength lidar observations in Northern France, we show systematically distinct microphysical properties of UTLS smoke aerosols compared to their free-tropospheric counterparts.

The analysis of satellite data reveals multiple episodes of smoke intrusions into the stratosphere through pyroconvection (PyroCb) and synoptic-scale processes (warm conveyor belt, WCB). Model simulations using MOCAGE chemistry-transport model, which included emission data from GFAS (Global Fire Assimilation System) are shown to accurately capture the synoptic-scale uplift of smoke into the UTLS and reproduce the spatial evolution of the aerosol plumes.

We show that the multiple episodes of wildfire-driven stratospheric intrusions during Boreal Summer 2023 through PyroCb and WCB mechanisms are altogether responsible for the recordhigh and persistent season-wide smoke pollution at the commercial aircraft cruising altitudes and the lowermost stratosphere.