



**HAL**  
open science

## Modeling soft X-ray emissions at the dayside magnetopause

Qiuyu Xu, Dimitra Koutroumpa, Ronan Modolo, Tianran Sun, Hyunju Connor, Steve Sembay

► **To cite this version:**

Qiuyu Xu, Dimitra Koutroumpa, Ronan Modolo, Tianran Sun, Hyunju Connor, et al.. Modeling soft X-ray emissions at the dayside magnetopause. EGU General Assembly 2024, Apr 2024, Vienna, Austria. pp. EGU24-1530, 10.5194/egusphere-egu24-1530 . insu-04515671

**HAL Id: insu-04515671**

**<https://insu.hal.science/insu-04515671>**

Submitted on 31 May 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EGU24-1530, updated on 31 May 2024

<https://doi.org/10.5194/egusphere-egu24-1530>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



## Modeling soft X-ray emissions at the dayside magnetopause

Qiuyu Xu<sup>1</sup>, Dimitra Koutroumpa<sup>1</sup>, Ronan Modolo<sup>1</sup>, Tianran Sun<sup>2</sup>, Hyunju Connor<sup>3</sup>, and Steve Sembay<sup>4</sup>

<sup>1</sup>LATMOS/IPSL, CNRS, UVSQ Université Paris-Saclay, Sorbonne Université, Guynacourt, France

<sup>2</sup>National Space Science Center, Chinese Academy of Sciences, Beijing, China

<sup>3</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>4</sup>Leicester University, Leicester, UK

In this study, we simulate the Solar Wind Charge Exchange (SWCX) soft X-ray emissions at dayside magnetopause by using magnetohydrodynamic (MHD) and Test-Particle (TP) models. Due to the single fluid description, the MHD model is unable to resolve the particle kinetic effects or distinguish the magnetospheric plasma and the solar wind plasma. We investigate these effects with the TP model. As the TP model does not self-compute magnetic and electric field, the magnetic and electric field data obtained from OpenGGCM and PPMLR MHD model are used as the input to TP model. The soft X-ray emissivity maps simulated from pure OpenGGCM and PPMLR MHD approaches and from TP-OpenGGCM and TP-PPMLR MHD approaches are presented and compared. The results indicate that the TP model can well resolve the kinetic effects and the individual spectral characteristics, and it does not require a masking method in the magnetosphere. Therefore the TP model is a complementary approach for simulating the X-ray emissions in the dayside magnetosheath and cusps.