



HAL
open science

Seasonal variation of trace species in Titan's ionosphere

Maélie Coutelier, Thomas Gautier, Koyena Das, Joseph Serigano, Sarah Horst

► **To cite this version:**

Maélie Coutelier, Thomas Gautier, Koyena Das, Joseph Serigano, Sarah Horst. Seasonal variation of trace species in Titan's ionosphere. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. pp.EPSC2022-479, 10.5194/epsc2022-479 . insu-04545510

HAL Id: insu-04545510

<https://insu.hal.science/insu-04545510>

Submitted on 14 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Europlanet Science Congress 2022

Palacio de Congresos de Granada, Spain

18 – 23 September 2022

EPSC Abstracts

Vol. 16, EPSC2022-479, 2022, updated on 23 Sep 2022

<https://doi.org/10.5194/epsc2022-479>

Europlanet Science Congress 2022

© Author(s) 2022. This work is distributed under
the Creative Commons Attribution 4.0 License.



Seasonal variation of trace species in Titan's ionosphere

Maélie Coutelier¹, Thomas Gautier¹, Koyena Das¹, Joseph Serigano², and Sarah Horst²

¹LATMOS, CNRS, Guyancourt, France (maelie.coutelier@latmos.ipsl.fr)

²Department of Earth and Planetary Sciences, Johns Hopkins University, Baltimore, MD, USA

Introduction

With 13 years of observations, the Ion and Neutral Mass Spectrometer (INMS) onboard the Cassini spacecraft has observed the upper atmosphere of Titan through two seasons: winter and spring. The complex atmosphere is mainly composed of N₂, CH₄, H₂ and Ar, but a lot more carbon and nitrogen bearing trace species have been observed by INMS and other instruments. Using data from the closed source neutral mode of INMS instrument, we studied the abundance and variation of traces neutral species in Titan ionosphere, between 1500 and 950 km of altitude. We will present an ongoing effort on the reanalysis of the entire INMS Titan's observation dataset.

Method

To do so we recalibrated INMS data by taking into account the dead time correction, the ram pressure enhancement, the saturation correction, the increase of pressure in the chamber with the decreases of altitude, the sensitivity and the contamination by thruster firing (Cui et al., 2009,2012). In addition, species entering the instrument were ionized and fragmented into ions inside INMS chamber, making difficult the identification of different species in such complex mass spectra. To retrieve the molecular mixing ratios we used a Monte-Carlo sampling on the fragmentation pattern to deconvolve the signal. To obtain a complete mass spectrum (m/z 1 to 99), we stacked INMS data, which increases the uncertainty on the altitude. We used the mass spectra deconvolution code developed by Gautier et al., (2020), also employed by Serigano et al., (2020) when they treated Saturn INMS data.

This enabled the retrieval of vertical and seasonal variation of Titan's atmosphere minor components. We expect to be able to link our results with the seasonal variations observed by other instruments [such as CIRS (Mathé et al., 2020)] in lower atmospheric layers.

References

Cui et al.(2009) Analysis of Titan's neutral upper atmosphere from Cassini Ion Neutral Mass Spectrometer measurements. *Icarus* 200 (2009) 581–615

Cui et al.(2012) The CH₄ structure in Titan's upper atmosphere revisited. *J. Geophys. Res.*, 117, E11006, doi:10.1029/2012JE004222.

Gautier et al. (2020) Decomposition of electron ionization mass spectra for space application using a Monte-Carlo approach. *Rapid. Com. Mass Spec.* 34(8), e8659

Mathé et al., (2020) Seasonal changes in the middle atmosphere of Titan from Cassini/CIRS observations: Temperature and trace species abundance profiles from 2004 to 2017. *Icarus*, 344, 113547.

Serigano et al. (2020) Compositional measurements of saturn's upper atmosphere and rings from cassini INMS. *Journal of Geophysical Research: Planets*, 125(8), e2020JE006427.

How to cite: Coutelier, M., Gautier, T., Das, K., Serigano, J., and Horst, S.: Seasonal variation of trace species in Titan's ionosphere, Europlanet Science Congress 2022, Granada, Spain, 18–23 Sep 2022, EPSC2022-479, <https://doi.org/10.5194/epsc2022-479>, 2022.



Discussion

[Login](#) to access the discussion