

The Vorticized Volcanic Plumean unusual stratospheric phenomenon revealed by Aeolus wind profiling

Sergey Khaykin, Jos de Laat, Sophie Godin-Beekmann, Alain Hauchecorne, Mathieu Ratynski

▶ To cite this version:

Sergey Khaykin, Jos de Laat, Sophie Godin-Beekmann, Alain Hauchecorne, Mathieu Ratynski. The Vorticized Volcanic Plumean unusual stratospheric phenomenon revealed by Aeolus wind profiling. Aeolus Science Conference 2023, May 2023, Rhodes Island, Greece. insu-04601342

HAL Id: insu-04601342 https://insu.hal.science/insu-04601342

Submitted on 4 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Vorticized Volcanic Plume – an unusual stratospheric phenomenon revealed by Aeolus wind profiling



Sergey Khaykin¹, Jos de Laat²,

Sophie Godin-Beekmann¹, Alain Hauchecorne¹, and Mathieu Ratynski¹




¹ Laboratoire Atmosphères, Observations Spatiales (LATMOS), INSU-CNRS, Guyancourt, France

² Royal Netherlands Meteorological Institute, De Bilt, the Netherlands












Raikoke eruption 21 June 2019





- Injection altitude: 14 17 km
- SO₂ injected mass: 1.4 2.1 Tg
- Fine ash injected mass: 0.4 1.8 Tg
- Strongest NH eruption since Pinatubo
- Unusual self-lofting of Raikoke plume rising diabatically by 11 km (up to 27 km)
- Behavior similar to wildfire smoke plumes




SCV – Smoke-Charged Vortex

Australian ANY fires 2020


PyroCb cloud Credits: @NWS Albuquerque

Compact bubble of smoke (CALIOP)




SCV – Smoke-Charged Vortex, confining the injected material and maintaining high concentration of absorptive aerosols inside the bubble and its heating

Khaykin et al., Nat. Com. EE, 2020




Any sizable injection of smoke into the stratosphere produces an SCV

Confinement of volcanic material (TROPOMI SO2)



Confinement of volcanic material




- Persistent compact and circular shape of the volcanic plume's core
- Bottom-side tailing and inverted comma shape
- Indications for anticyclonic motion but no clear confirmation from ERA5 data

Khaykin et al., Sci. Rep., 2022

VVP - Vorticized Volcanic Plume

Aeolus to provide unique observational evidence for rotation of the plume





VVP in ERA5




ERA5 Potential vorticity












ERA5 reproduces the PV anomaly associated with the plume

Fairbanks, 65 N 148W: 7-8 Jul 2019




Internal heating of the plume





Warm anomaly:

- 1 K (GNSS-RO)
- 0.6 K (ERA5)
- Up to 2 K (RS)



32

34

Longitude (deg.)


36


-0.5

Latitude (deg)


Self-lofting of Raikoke VVP

Diabatic ascent at 2.7 K/day rate up to 27 km






VVP tracking for 3+ months, 3+ circumnavigations





Acceleration of diabatic rise during ASM overpasses


Raikoke vs Sarychev

- Raikoke eruption was rich in fine ash (strong and persistent depolarization of the plume) as opposed to Sarychev eruption
- No self-lofting of Sarychev plume
- Prolonged residence time of Raikoke aerosol in the stratosphere












Confined fraction of Raikoke SO2 emissions




 25% of the erupted SO₂ mass was contained by the VVP and was subject to significant self-lofting and equatorward transport




Aeolus observations of Australian Smoke-Charged Vortex






SCV dynamical structure from lidars and ERA5







SCV transit over Lauder, NZ (color map = depolarization, contous = Vwind summer anomaly)



Date in February 2020

CALIOP transects of SCV near New Zealand





3D structure of the vortex from ERA5 PV

Summary

Volcanic plumes are capable of self-organizing into persistent anticyclonic vortices under certain conditions

Required ingredients for VVP/SCV formation:

- Absorbing aerosols (ash, smoke)
- Extratropical location (PV gradient across the tropopause)
- Summer season (calm stratosphere)
- Dynamical confinement, maintaining the absorptive aerosols at high concentration is the key prerequisite for self-lofting
- Self-lofting of wildfire and volcanic plumes alters their global transport and prolongs their atmospheric residence time and radiative effects
- ESA frontier missions provide new insights into dynamics of volcanic and wildfire plumes

nature > scientific reports > articles > article

Article Open Access | Published: 27 December 2022

Unexpected self-lofting and dynamical confinement of volcanic plumes: the Raikoke 2019 case


S. M. Khaykin , A. T. J. de Laat , S. Godin-Beekmann, A. Hauchecorne & M. Ratynski




News / ESA's wind mission helps inves.


ESA's wind mission helps investigate the nature of volcanic plumes

Summary





VVP - Vorticized Volcanic Plume



VVP overpass above Hawaii

