



**HAL**  
open science

## **Constraining ion transport in the diamagnetic cavity of comet 67P**

Zoe Lewis, Arnaud Beth, Marina Galand, Pierre Henri, Martin Rubin, Peter Stephenson

### ► **To cite this version:**

Zoe Lewis, Arnaud Beth, Marina Galand, Pierre Henri, Martin Rubin, et al.. Constraining ion transport in the diamagnetic cavity of comet 67P. European Geosciences Union General Assembly 2024 (EGU24), Apr 2024, Vienne, Austria. pp.5481, <10.5194/egusphere-egu24-5481>. <insu-04652307>

**HAL Id: insu-04652307**

**<https://insu.hal.science/insu-04652307v1>**

Submitted on 18 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

EGU24-5481, updated on 18 Jul 2024

<https://doi.org/10.5194/egusphere-egu24-5481>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



## Constraining ion transport in the diamagnetic cavity of comet 67P

Zoe Lewis<sup>1</sup>, Arnaud Beth<sup>1</sup>, Marina Galand<sup>1</sup>, Pierre Henri<sup>2,3</sup>, Martin Rubin<sup>4</sup>, and Peter Stephenson<sup>5</sup>

<sup>1</sup>Imperial College London, Department of Physics, United Kingdom of Great Britain – England, Scotland, Wales

(z.lewis21@imperial.ac.uk)

<sup>2</sup>Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS, Université d'Orléans, Orléans, France (pierre.henri@cnrs-orleans.fr)

<sup>3</sup>Laboratoire Lagrange, OCA, UCA, CNRS, Nice, France (pierre.henri@cnrs-orleans.fr)

<sup>4</sup>Physikalisches Institut, University of Bern, Bern, Switzerland (martin.rubin@unibe.ch)

<sup>5</sup>Lunar and Planetary Laboratory, University of Arizona, USA (pstephenson@arizona.edu)

Comets are small icy bodies originating from the outer solar system that produce an increasingly dense gas coma through sublimation as they approach perihelion. Photoionisation of this gas results in a cometary ionosphere, which interacts with the impinging solar wind, leading to large scale plasma structures. One such structure is the diamagnetic cavity: the magnetic field-free inner region that the solar wind cannot penetrate. This region was encountered many times by the ESA Rosetta mission, which escorted comet 67P/Churyumov-Gerasimenko for a two-year section of its orbit.

Within the diamagnetic cavity, high ion bulk velocities have been observed by the Rosetta Plasma Consortium (RPC) instruments. The fast ions are thought to have been accelerated by an ambipolar electric field, but the nature and strength of this field are difficult to determine analytically. Our study therefore aims to model the impact of various electric field profiles on the ionospheric density profile and ion composition. The 1D numerical model we have developed includes three key ion species (H<sub>2</sub>O<sup>+</sup>, H<sub>3</sub>O<sup>+</sup>, and NH<sub>4</sub><sup>+</sup>) in order to assess the sensitivity of each to the timescale of plasma loss through transport. NH<sub>4</sub><sup>+</sup> is of particular interest, as it has been previously shown to be the dominant ion species at low cometocentric distances near perihelion. It is only produced through the protonation of NH<sub>3</sub>, a minor component of the neutral gas, and we show that this makes it particularly sensitive to the electric field.

We also compare the simulated electron density to RPC datasets, to find the electric field strength and profile which best recreate the plasma densities measured inside the diamagnetic cavity near perihelion. From this, we also constrain the radial bulk ion speed that is required to explain the observations with the model.