



**HAL**  
open science

## Mesoscale Modelling of Titan's Shangri-La region

Maxence Lefevre, Léa Bonnefoy, Aymeric Spiga

► **To cite this version:**

Maxence Lefevre, Léa Bonnefoy, Aymeric Spiga. Mesoscale Modelling of Titan's Shangri-La region. Europlanet Science Congress 2024, Sep 2024, Berlin, Germany. pp.EPSC2024-408, 10.5194/epsc2024-408 . insu-04676937

**HAL Id: insu-04676937**

**<https://insu.hal.science/insu-04676937v1>**

Submitted on 24 Aug 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Mesoscale Modelling of Titan's Shangri-La region

Maxence Lefevre<sup>1</sup>, Léa Bonnefoy<sup>2</sup>, and Aymeric spiga<sup>2</sup>

<sup>1</sup>Sorbonne Université, LATMOS, Paris, France (maxence.lefevre@latmos.ipsl.fr)

<sup>2</sup>Sorbonne Université, LMD, Paris, France

### 1. Introduction

Titan's weather is particularly active at all spatial scales: global (e.g., super-rotating winds), intermediate / mesoscale (e.g., convective thunderstorms, air-sea circulations), local (e.g., turbulence in the planetary boundary layer, the atmospheric layer in direct contact with the surface). The aeolian environment of Titan is active, with a fifth of Titan's surface being covered by dunes fields and the organic dust particles at the surface being potentially transported by wind circulations at all scales.

In 2034 NASA's Dragonfly quadcopter will land and explore Selk impact crater structure in the dark Shangri-La region [1]. This area is characterized by a surface rich in organic dust particles organized as dune fields, whose peculiar morphology probably involves two distinct wind regimes that remain to be understood [2].

### 2. Model

To study the near-surface dynamics, a new model was developed based on the Weather-Research Forecast (WRF) non-hydrostatic dynamical core [3] and coupled with the LMD Titan PCM physics package [4]. The domain of interest is centered on Selk crater and Shangri-La region. The horizontal resolution is set to 5 km. High-resolution topography and surface characteristic such as albedo, thermal inertia and surface roughness, were created based on Cassini SAR images. The detail of these synthetic maps are described in detail in the poster presented in this session by Bonnefoy et al. The meteorological fields initial states are taken from LMD Titan PCM outputs. The mesoscale boundary conditions are forced with the LMD Titan GCM.

### 3. Results

Fig 1 shows Snapshots map of horizontal winds 10 m above local surface (m/s) at noon. The topography, plains and mountains, affects the direction of the surface winds. The vast majority of the wind is between  $\pm 0.5$  m/s, consistent with observations. The high mountains are able to engender mountain waves, with horizontal wind reaching 1.0 m/s. The diurnal cycle has an impact on the amplitude and direction of wind, with preferably anabatic winds during the day and katabatic winds at night. The albedo, thermal inertia and surface roughness are set constant respectively to 0.2, 335 J K<sup>-1</sup> m<sup>-3</sup> and 5 cm. The impact of the spatial variability due to terrain type of these parameters on the winds will be discussed.




Figure 1: Snapshots map of horizontal winds 10 m above local surface ( $\text{m s}^{-1}$ ) at noon in the center of the domain. The red circle represent the landing site of Dragonfly [1].

#### References

- [1] Lorenz, R. D. et al. Selection and Characteristics of the Dragonfly Landing Site near Selk Crater, Titan. *Planet. Sci. J.* 2, 24 (2021).
- [2] Malaska, M. J. et al. Geomorphological map of the Afekan Crater region, Titan: Terrain relationships in the equatorial and mid-latitude regions. *Icarus* 270, 130–161 (2016).
- [3] Skamarock, W. C. & Klemp, J. B. A time-split nonhydrostatic atmospheric model for weather research and forecasting applications. *Journal of Computational Physics* 227, 3465–3485 (2008).
- [4] Lebonnois, S., Burgalat, J., Rannou, P. & Charnay, B. Titan global climate model: A new 3-dimensional version of the IPSL Titan GCM. *Icarus* 218, 707–722 (2012).