



Decline in Water Ice Abundance in the Martian Mesosphere during Aphelion

Daniel Toledo, Pascal Rannou, Victor Apestigue, Raul Rodriguez-Veloso, Ignacio Arruego, German Martinez, Leslie Tamppari, Asier Munguira, Ralph Lorenz, Aurelien Stcherbinine, et al.

► To cite this version:

Daniel Toledo, Pascal Rannou, Victor Apestigue, Raul Rodriguez-Veloso, Ignacio Arruego, et al.. Decline in Water Ice Abundance in the Martian Mesosphere during Aphelion. Europlanet Science Congress 2024, Europlanet Science Congress 2024, Sep 2024, Berlin, Germany. pp.PSC2024-504, 10.5194/epsc2024-504 . insu-04685335

HAL Id: insu-04685335

<https://insu.hal.science/insu-04685335v1>

Submitted on 3 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EPSC Abstracts

Vol. 17, EPSC2024-504, 2024, updated on 03 Sep 2024

<https://doi.org/10.5194/epsc2024-504>

Europlanet Science Congress 2024

© Author(s) 2024. This work is distributed under
the Creative Commons Attribution 4.0 License.



Decline in Water Ice Abundance in the Martian Mesosphere during Aphelion

Daniel Toledo¹, Pascal Rannou², Victor Pestigue¹, Raul Rodriguez-Veloso¹, Ignacio Arruego¹, German Martinez³, Leslie Tamppari⁴, Asier Munguira⁵, Ralph Lorenz⁶, Aurelien Stcherbinine⁷, Franck Montmessin⁸, Agustin Sanchez-Lavega⁵, Priya Patel⁴, Michael Smith⁹, Mark Lemmon¹⁰, Alvaro Vicente-Retortillo¹¹, Claire Newman¹², Daniel Viudez-Moreiras¹¹, Ricardo Hueso⁵, Tanguy Bertrand¹³, and the +4*

¹Instituto Nacional de Técnica Aeroespacial (INTA), Madrid, Spain

²GSMA, UMR 7331-GSMA, Université de Reims Champagne-Ardenne, 51687 Reims, France

³Lunar and Planetary Institute, Universities Space Research Association, Houston, TX, USA

⁴Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

⁵Universidad del País Vasco UPV/EHU, Bilbao, Spain

⁶Johns Hopkins Applied Physics Laboratory, Laurel, MD, USA

⁷IRAP - Institut de Recherche en Astrophysique et Planétologie, Toulouse, France

⁸Laboratoire Atmosphères Milieux Observations Spatiales (LATMOS), Université Paris-Saclay, Sorbonne Université, Centre

National de la Recherche Scientifique, Guyancourt, France

⁹NASA Goddard Space Flight Center, Greenbelt, MD, USA

¹⁰Space Science Institute, Boulder, CO, USA

¹¹Centro de Astrobiología (INTA-CSIC), Torrejón de Ardoz, Madrid, Spain

¹²Aeolis Research, Chandler, AZ, USA

¹³LESIA, Observatoire de Paris, Meudon, France

*A full list of authors appears at the end of the abstract

Clouds play a crucial role in the past and current climate of Mars. Cloud particles impact the planet's energy balance and atmospheric dynamics, as well as influence the vertical distribution of dust particles through dust scavenging. This process of dust scavenging by clouds has significant consequences for the planet's water cycle. For example, regions in the atmosphere with insufficient quantities of dust particles, or condensation nuclei, can inhibit the formation of H₂O clouds, leading to the presence of water vapor in excess of saturation [1]. Recent observations made by the MEDA Radiation and Dust Sensor (RDS) [2,3] have shown a marked decline in mesospheric cloud activity (above 35-40 km) when Mars is near its aphelion (within the Aphelion Cloud Belt-ACB season), notably occurring during solar longitudes (Ls) between Ls 70° and 80° [4] (see Figure 1).

In order to investigate the possible factors leading to this decrease in water ice abundance, we used a one-dimensional cloud microphysical model [5,6], which includes the processes of nucleation, condensation, coagulation, evaporation, precipitation, and coalescence, and where the vertical mixing is parameterized using an eddy diffusion profile (K_{eddy}). Combining cloud microphysics modeling with ground-based (Mars 2020 and InSight) and orbital observations (TGO and MRO) of clouds, water vapor, and temperature, we will discuss in this presentation the main factors controlling the water abundance in the Martian mesosphere during the ACB season.

References: [1] Maltagliati, Luca, et al. "Evidence of water vapor in excess of saturation in the atmosphere of Mars." *science* 333.6051 (2011): 1868-1871. [2] Apestigue, V., et al. "Radiation and Dust Sensor for Mars Environmental Dynamic Analyzer Onboard M2020 Rover". *Sensor* 22.8 (2022): 2907. [3] Rodriguez-Manfredi, Jose Antonio, et al. "The Mars Environmental Dynamics Analyzer, MEDA. A suite of environmental sensors for the Mars 2020 mission." *Space science reviews* 217.3 (2021): 1-86. [4] Toledo, D., et al. "Measurement of aerosol optical depth and sub-visual cloud detection using the optical depth sensor (ODS)". *Atmospheric Measurement Techniques* 9.2 (2016): 455-467. [5] Montmessin, F., Rannou, P., Cabane, M.: New insights into martian dust distribution and water-ice cloud microphysics. *Journal of Geophysical Research: Planets* 107(E6), 41 (2002). [6] Rannou, P., Montmessin, F., Hourdin, F., Lebonnois, S.: The latitudinal distribution of clouds on titan. *science* 311(5758), 201205 (2006).




Fig 1. Mesospheric cloud opacities retrieved from MEDA-RDS observations at twilight for Martian years (MY) 36 and 37. In most of the cloud detections, the cloud altitudes were between 40 and 50 km.

+4: Jorge Pla-Garcia¹¹, Margarita Yela¹, Manuel de la Torre Juarez⁴, Jose Antonio Rodriguez-Manfredi¹¹