



**HAL**  
open science

## Editorial: Volcanic Islands-A Challenge for Volcanology

Alessandro Bonforte, Joan Martí, Antonio Paonita, Michel Pichavant

► **To cite this version:**

Alessandro Bonforte, Joan Martí, Antonio Paonita, Michel Pichavant. Editorial: Volcanic Islands-A Challenge for Volcanology. *Frontiers in Earth Science*, 2022, 10, 10.3389/feart.2022.954902 . insu-04702181

**HAL Id: insu-04702181**

**<https://insu.hal.science/insu-04702181v1>**

Submitted on 19 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Editorial: Volcanic Islands—A Challenge for Volcanology

Alessandro Bonforte<sup>1\*</sup>, Joan Martí<sup>2</sup>, Antonio Paonita<sup>3</sup> and Michel Pichavant<sup>4</sup>

<sup>1</sup>Istituto Nazionale di Geofisica e Vulcanologia (INGV), Sezione di Catania - Osservatorio Etno, Catania, Italy, <sup>2</sup>Geosciences Barcelona, CSIC, Barcelona, Spain, <sup>3</sup>Istituto Nazionale di Geofisica e Vulcanologia (INGV), Sezione di Palermo - Osservatorio Etno, Palermo, Italy, <sup>4</sup>Institut des Sciences de la Terre d'Orléans, CNRS, Orléans, France

**Keywords:** eruption, magma-water interaction, seafloor, submarine volcanic edifice, eruptive stratigraphy, eruptive scenarios, hazard and risk

## Editorial on the Research Topic

### Volcanic Islands - A Challenge for Volcanology

Most volcanoes on the Earth rise from the bottom of seas and oceans. Most of them do not reach the surface of sea and remain hidden to all conventional observations from surface and space. Only some of them rise above the sea level, forming islands and passing from submarine to subaerial volcanism. Volcanic islands develop in virtually all the geodynamic contexts on Earth, from mid-ocean ridges (Iceland), to intraplate (Hawaii), to volcanic arcs (Aeolian Islands). All the liquid-descent evolutive degrees of magma are finally represented, from primitive compositions up to strongly evolved rhyolite, trachyte and phonolite lavas. So, the eruptive styles of these volcanoes range consequently from mild effusions to plinian eruptions.

The interface with the sea poses some particular conditions due to the interaction with seawater, both at surface and below. Large hydrothermal activity easily develops due to the circulation of seawater at depth, and consequent interaction with magmatic gases and hot country rocks. The presence of the water and sea erosion also triggers particular instability conditions of the slopes of these volcanic edifices, so that the effects by seismicity and ground deformations can become disastrous. Volcanic islands are by far the most vulnerable environments on Earth, not only because of their high exposure to the impact of multi-hazards, where several hazardous phenomena may interact in a simultaneous or consecutive way, but also due to their isolation from the main land, which makes external supply chains and help during a crisis difficult. A large number of volcanic islands are in fact intensively inhabited, despite the limited available lands, thus the common volcanic phenomenologies pose severe hazardous scenarios and submarine eruptions and landslides triggering catastrophic tsunami waves have to be considered.

One of the most important Research Topics to be considered when investigating and monitoring island volcanoes, is that most of the edifice is submerged and the emerging part only constitutes the summit of the volcano. This poses enormous difficulties to scientists, both from a scientific and technological point of view, because the development of monitoring systems at seafloor is still in its infancy. On this ground, such edifices represent one of the most important and interesting challenges for volcanologists and Earth scientists in general.

This Research Topic collected 10 contributions, covering a very wide range of disciplines, from petrology to submarine geophysics, all aimed at understanding and characterizing the activity, dynamics and hazards of volcanoes developing at the interface between land and sea, not only forming islands but also raising above the sea along a continental coastline. This latter is the case of Mt. Etna, whose submerged eastern flank has been investigated in one of the contributions (Urlaub et al.), describing and analyzing a wide series of geophysical imaging and seafloor geodetic surveys

## OPEN ACCESS

### Edited and reviewed by:

Valerio Acocella,  
Roma Tre University, Italy

### \*Correspondence:

Alessandro Bonforte  
alessandro.bonforte@ingv.it

### Specialty section:

This article was submitted to  
Volcanology,  
a section of the journal  
Frontiers in Earth Science

Received: 27 May 2022

Accepted: 07 June 2022

Published: 22 June 2022

### Citation:

Bonforte A, Martí J, Paonita A and  
Pichavant M (2022) Editorial: Volcanic  
Islands—A Challenge for Volcanology.  
Front. Earth Sci. 10:954902.  
doi: 10.3389/feart.2022.954902

that unveil some volcano-tectonic structures probably representing the link between flank dynamics and tectonic triggering of the instability.

Arc volcanic islands have also been investigated in this Research Topic, that is the case of the Aleutian Arc segment (Fischer et al.). Geochemical data show that all these volcanoes host high-temperature magmatic-hydrothermal systems and have gas discharges typical of volcanoes in oceanic arcs. Volcanic CO<sub>2</sub> fluxes from this arc segment are small, probably due to the lower sediment flux delivered to the trench in this part of the arc.

In volcanic islands, stratigraphy and sedimentary records can be particularly useful to reconstruct past volcanic activity. One of the contributions (Andrade et al.) focused on Flores island (Azores) shows in fact that the volcano experienced vigorous volcanic activity during the Late Holocene; therefore, contrary to what is assumed, the possibility of future eruptions should be properly considered, and the volcanic hazard here should not be underestimated.

The Hazard aspects have been investigated also for remote islands, in a contribution (Gjerløw et al.) focused on Jan Mayen Island, in the North-Atlantic. The eruptive history of this island, showing variable activity styles, has been exploited for evaluating the different vent opening probabilities, exposure to different volcanic phenomena and recurrence times for depicting different scenarios and qualitative hazard maps.

A similar hazard evaluation come from the PDCs deposits on Terceira island (Pimentel et al.). Studies on the geometry and structure of two ignimbrite deposits on the island reveal interesting details on the eruptions that produced them. Even if ignimbrite-forming eruptions are not frequent on that volcano, this kind of hazard is significant, since its effect can be dramatic for the population.

Geochronological and isotopes analyses, as well as field surveys and petrological analysis were used for reconstructing the eruptive history of Weizhou island in the Guangxi Province (China), for evaluating the risk to inhabitants (Yu et al.). A very variable eruptive style has been found on the four volcanoes discovered, spanning from explosive, PDC-forming eruptions to effusive ones, also with some evidences of magma-water interaction.

The Canary Islands are a classical site locality for alkaline, SiO<sub>2</sub>-undersaturated intraplate magmas (Klügel et al.). Products from one eruption of the now active Cumbre Vieja volcano of La Palma contain abundant inclusions from different levels of the plumbing system, mantle rocks, oceanic gabbros and cumulates.

Phonolites form beneath oceanic island volcanoes by combined differentiation and periodic recharge by mantle magmas. They require a balance between rates and volumes of magma recharge pulses and of eruptive events.

In small islands such as Martinique (Lesser Antilles), the history of the volcanic activity is generally better preserved in submarine deposits than on-land (Villemant et al.). Cores of marine sediments reconstitute the chronostratigraphy of tephra, volcanoclastic turbidites, and mass-wasting events over 1.5 Ma. Results provide a framework for further petrological studies of Mt Pelée and Pitons du Carbet and for the timing of flank collapses and submarine landslides.

The very recent eruptive history of a newborn and growing island has been analyzed and correlated with the variations of the chemical composition of erupted magma (Maeno et al.). Results of this analysis have been used to infer the minimum eruption rates and durations required for forming volcanic islands able to survive to the erosion.

Volcanism on ocean island can also serve as windows into the composition and behavior of mantle plumes (Wilson et al.). Santa Cruz volcano (Galapagos) followed a distinct geochemical evolution from the present-day western shields, having been constructed under the influence of the Galápagos Spreading Center. Ocean island evolution can change dramatically depending on the plume's tectonic setting.

## AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

*Copyright © 2022 Bonforte, Martí, Paonita and Pichavant. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.*