



HAL
open science

High stabilization of pentavalent uranium on magnetite nanoparticles evidenced by high-energy-resolution X-ray absorption spectroscopy

Takumi Yomogida, Jaimy Scaria, Laura Fablet, Kohei Tokunaga, Syuntaro Dei, Kotaro Higashi, Naomi Kawamura, Yoshio Takahashi, Rémi Marsac

► To cite this version:

Takumi Yomogida, Jaimy Scaria, Laura Fablet, Kohei Tokunaga, Syuntaro Dei, et al.. High stabilization of pentavalent uranium on magnetite nanoparticles evidenced by high-energy-resolution X-ray absorption spectroscopy. *Chemical Communications*, 2025, 61 (91), pp.17926-17929. <10.1039/D5CC04540A.>. <insu-05314225>

HAL Id: insu-05314225

<https://insu.hal.science/insu-05314225v1>

Submitted on 14 Oct 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. Yomogida, J. Scaria, L. Fablet, K. TOKUNAGA, S. Dei, K. Higashi, N. Kawamura, Y. Takahashi and R. Marsac, *Chem. Commun.*, 2025, DOI: 10.1039/D5CC04540A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

High stabilization of pentavalent uranium on magnetite nanoparticles evidenced by high-energy-resolution X-ray absorption spectroscopyReceived 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Takumi Yomogida,^{*a,b,c} Jaimy Scaria^d, Laura Fablet^d, Kohei Tokunaga^e, Syuntaro Dei^f, Kotaro Higashi^g, Naomi Kawamura^g, Yoshio Takahashi^c and Rémi Marsac^{*b}**We observed that U(V) is more stable on magnetite nanoparticles than expected, across a wide range of pH and redox conditions and even after 10 days of reaction, as revealed by high-energy-resolution fluorescence-detection X-ray absorption spectroscopy combined with multivariate curve resolution.**

Uranium(U), widely used as a fuel for nuclear power generation, is a key element in actinide chemistry, nuclear technology, and environmental science due to its usefulness as an energy resource and its chemical toxicity¹⁻³. In the environment, it has traditionally been considered to exist as U(IV) and U(VI). Although long overlooked^{4,5}, recent studies have highlighted the significance of U(V), notably when U(VI) interacts with Fe(II)-bearing minerals, such as magnetite^{6,7-11}. Magnetite is a ubiquitous mineral in natural settings and may also form in artificial barriers for nuclear waste disposal as a result of steel corrosion¹². However, the role of U(V) remains unresolved as it is either considered as a metastable reaction intermediate during U(VI) reduction to U(IV) at magnetite surfaces⁹ or as a stable species when incorporated via coprecipitation with magnetite¹¹. As a result, the retention of U by magnetite has been intensively studied in recent years^{9,11,13-15}. Nonetheless, the mechanisms controlling U interaction with

magnetite in aqueous solution relevant to natural systems remain poorly understood. In particular, many studies have employed high U concentrations, in which the precipitation of UO_{2(s)} nanoparticles is often observed. In contrast, single-atom surface species formation may be a more relevant mechanism to investigate in many environments where U occurs at trace levels⁵, as was also recently shown for rhenium.¹⁶

Recently, high-energy-resolution fluorescence detection (HERFD)-XANES has been applied to a variety of U compounds as a more sensitive technique for analysing electronic structures¹⁷⁻²⁷. The spectral features of HERFD-XANES are more distinct than those of conventional XANES spectra. For example, our group observed a specific peak splitting of U(V) in FeUO₄ with U L₃-edge HERFD-XANES, due to the strong crystal field splitting of the 6d orbital, which could not be detected in normal XANES spectra.²⁸ This specific peak splitting is not observed in HERFD-XANES spectra of U(IV) in UO₂ because of its weaker crystal-field splitting than that of U(V) in FeUO₄. Therefore, these promising results suggest that HERFD-XANES spectra could be a powerful technique to unravel the role played by U(V) when interacting with Fe-bearing minerals such as magnetite nanoparticles.

This paper presents insights into the influence of magnetite stoichiometry ($0 \leq R = \text{Fe(II)/Fe(III)} \leq 0.5$) on the surface reduction of U(VI) to U(V) and U(IV), as a key parameter controlling U redox speciation in natural settings. Although R can readily change due to the oxidation of structural Fe(II) or proton/ligand-promoted dissolution²⁹⁻³¹, prior studies have not quantified U(V) when assessing these effects^{4,5}. To address this gap, we employed U L₃-edge HERFD-XANES spectroscopy to investigate the electronic structure of U on magnetite with varying stoichiometries and observed a peak splitting in the HERFD-XANES spectrum of U on magnetite, an effect not previously detected using conventional XANES. In this study, we explore whether U(V) stabilization correlates with structural Fe(II) in magnetite by comparing U reference materials of U(IV)O₂, FeU(V)O₄, and U(VI)O₂(NO₃)₂(H₂O)₆, for which HERFD-XANES analyses have already been reported²⁸.

^a Nuclear Safety Research Center, Japan Atomic Energy Agency, 2-4 Shirakata, Naka, Tokai, Ibaraki, 319-1195, Japan^b Université Paris Cité, Institut de physique du globe de Paris, CNRS, F-75005 Paris, France.^c Department of Earth and Planetary Science, The University of Tokyo, Hongo 7-3-1, Bunkyo, Tokyo 113-0033, Japan^d Univ Rennes, CNRS, Géosciences Rennes – UMR 6118, F-35000 Rennes, France^e Ningyo-toge Environmental Engineering Center, Japan Atomic Energy Agency, Tomata-gun, Okayama, 708-0601, Japan^f Horonobe Underground Research Center, Japan Atomic Energy Agency, 432-4 Hokushin, Horonobe, Hokkaido 098-3224, Japan^g Center for Synchrotron Radiation Research, Japan Synchrotron Radiation Research Institute (JASRI), Sayo, Hyogo 679-5198, Japan

*Corresponding authors

E-mail: yomogida.takumi@jaea.go.jp

†Electronic Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

COMMUNICATION

ChemComm

Stock suspensions of 10 nm-sized stoichiometric ($R = 0.5$) and non-stoichiometric ($R = 0.1, 0.2, 0.3$ and 0.4) magnetite were synthesized and characterized as described in previous studies^{31,32}. The U stock solution (0.5 mM) was prepared from uranyl acetate and dissolved in 0.01 M HCl solution. The uranium concentration (10 μM , 2.3 ppm), background electrolyte concentration (10 mM NaCl), and total Fe concentration in the suspension (20 mM; equivalent to 1.1 g L⁻¹ for stoichiometric magnetite) were set equal for all experiments. As the surface area of 10 nm-sized magnetite nanoparticles is approximately 100 m² g⁻¹, the maximum surface loading corresponds to 0.1 $\mu\text{mol m}^{-2}$ (0.06 atom nm⁻²), which ensures that surface complexation was the predominant mechanism over U precipitation⁵. Batch experiments were conducted in 50 mL polypropylene tubes. Dilute HCl or NaOH solutions were used to adjust the solution pH. Adsorption and reduction experiments of U(VI) were performed (i) for 6 distinct pH values (4, 5, 6, 7, 8, and 9) onto stoichiometric magnetite (R0.5) and (ii) at pH 6 and 8 onto non-stoichiometric magnetite (R0.3 and R0.2) at room temperature under anaerobic condition (N₂ - glovebox, JACOMEX, O₂ < 10 ppm). In addition, interaction of U(VI) with R0.1 was studied under ambient atmospheric conditions (i.e. in presence of O_{2(g)}); a reference system in which U(VI) is expected stable. After 10 days of reaction time, the solid phase was separated from the liquid phase by magnetic separation and dried in the glovebox. The dried samples were triple sealed using 100 μm high-barrier film (AP-1522, ISO). The HERFD-XANES experiments were performed at beamline BL39XU, SPring-8, as previously reported²⁸. A combined energy resolution of ~ 1.5 eV was achieved in the experiments, as determined by measuring the full width at half-maximum (FWHM). Data processing was carried out using the Fastosh software³³ and Athena software³⁴. Detailed experimental conditions are described in the Supplementary Information (SI).

The HERFD-XANES spectra at the U L₃ edge of U interacting with R0.5 at pH 6 (denoted R0.5pH6), collected by monitoring the L α_1 emission line, and the corresponding XANES spectrum collected in the conventional fluorescence mode are compared in Fig. 1a. The white line at approximately 17176 eV in the conventional XANES spectrum of R0.5pH6 mainly originates from the electron transition from U 2p_{3/2} to unoccupied 6d states. The HERFD-XANES spectrum of the same sample, the red line in Fig. 1a, revealed more distinct features: the white line was split into two peaks in the HERFD-XANES spectra. The HERFD-XANES spectra at the U L₃ edge of R0.5 pH6, UO₂, FeUO₄, and UO₂(NO₃)₂(H₂O)₆ are compared in Fig. 1b. Peak splitting was observed in the HERFD-XANES spectrum of FeUO₄, whereas no peak splitting was observed in the HERFD-XANES spectra of UO₂ and UO₂(NO₃)₂(H₂O)₆, indicating that a significant fraction of U is stabilized as U(V) species on magnetite.

Figure 2a shows the HERFD-XANES spectra of U-R0.5 samples prepared at different pH values. Only one peak was observed at 17175 eV at pH 4. As the pH increased, the intensity of the first peak decreased, while the second peak around 17180 eV increased. Figures 2b and 2c show the HERFD-XANES spectra of U-magnetite samples with different stoichiometries at pH 6 and 8, respectively. HERFD-XANES analysis revealed a




Figure 1. (a) U L₃ edge HERFD-XANES spectrum (red line) and XANES spectrum (black line) of U-magnetite samples prepared at pH 6. (b) (Solid line) U L₃ edge HERFD-XANES spectrum of U-magnetite samples prepared at pH 6. (Dotted line) HERFD-XANES spectra of UO₂(black), FeUO₄ (red), and UO₂(NO₃)₂ · 6H₂O(blue), previously published by Yomogida et al.




Figure 2. U L₃-edge HERFD-XANES spectra of U reacted with (a) stoichiometric magnetite (R=0.5) nanoparticles at different pH, (b) magnetites with different stoichiometries at pH 6 and (c) pH 8

gradual decrease in the second peak intensity (17180 eV) with decreasing stoichiometry. This trend was observed at both pH 6 (Fig. 2b) and pH8 (Fig. 2c). These observations confirm previous observations that the U redox speciation is influenced by R, with higher extent of U(VI) reduction in Fe(II)-rich system⁴. The effect of pH, which jointly affects U redox speciation in solution and R with the redox potential (E_h),³⁰ will be discussed in more detail later in this manuscript.

To quantify the number of U species on stoichiometric and non-stoichiometric magnetite, singular value decomposition (SVD) analysis was applied to HERFD-XANES spectra of U on magnetites using FASTOSH software³³. Figure S1 shows the dependence of the singular values on the number of components (n), obtained by the SVD of the data matrices. Because the singular values were nearly zero for $n > 4$, we considered that 3 components were sufficient to analyze the experimental data. Figures 3a, 3b, and 3c show the pure XANES spectra of components 1 – 3, obtained by multivariate curve resolution with alternating least squares (MCR-ALS) analysis of the data³⁵. Figure 3a shows the XANES of component 1, which displays a broad white line peak. The features of the XANES of the component 1 are consistent with those of the UO₂. Figure 3b shows the XANES the component 2, whose main peak is split and agrees with that of FeUO₄. Figure 3c shows the XANES spectrum of component 3, whose main peak is the narrowest




Figure 3. Pure XANES spectra of (a) component 1, (b) component 2, and (c) component 3 in MCR-ALS analysis of the data matrix using the PureS method. (Dotted line) HERFD-XANES spectra of UO_2 (black), FeUO_4 (red), and $\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (blue), as previously published in Yomogida et al. Percentage of fractions of U(IV), U(V), and U(VI) versus (d) different pH, (e) different magnetite stoichiometry at pH 6, and (f) different magnetite stoichiometry at pH 8

but agrees relatively well with that of $\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$. Thus, we assign the U surface species on magnetite to the 3 relevant oxidation states of U.

Figure 3d, e, and f show the distribution (%) of the 3 components obtained by MCR-ALS. Linear combination fitting (LCF) of XANES spectra of U on magnetite, using reference spectra of UO_2 , FeUO_4 and $\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$, also provided results similar to MCR-ALS (Figure S2). On stoichiometric magnetite, approximately half of the U bound to magnetite at $\text{pH} > 6$ was U(V) (Figure 3d). Thermodynamic speciation calculations in solution may support this observation, E_H values determined in R0.5 suspensions³⁰ approach the predominant field of U(V) for $\text{pH} 5.5\text{--}7.5$ (Figure S3). However, such calculations do not account for the formation of surface species, which generally favour tetra- and hexavalent oxidation states of actinides ($\text{An} = \text{U}, \text{Np}, \text{Pu}$) over the pentavalent ones. Indeed, actinides(V) weakly sorb to iron oxide surfaces because of the smaller effective charge of cation ($\text{U(V)O}_2^+ < \text{U(VI)O}_2^{2+} < \text{U}^{4+}$)^{36,37}. In addition, for $\text{pH} > 7.5$, the theoretical prevalence of aqueous U(IV) contrasts with the observed prevalence of U(V) at the magnetite surface. Therefore, this unexpectedly high stability of U(V) is difficult to explain solely by surface complexation processes. This may instead result from its favorable incorporation into the magnetite structure. Indeed, spectroscopic studies evidenced that U(V) was highly stable when coprecipitated with Fe during magnetite synthesis⁶, or that other ions such as Ni(II) ³⁸, Co(II) ³⁹, or Cr(III) ³² incorporated into magnetite during sorption experiments. The XANES spectra

of U(V) (Fig. 3b) is almost consistent with U(V) in FeUO_4 . U(V) may partially substitute octahedral site in magnetite. At $\text{pH} < 6$, magnetite can partially reduce U(VI) to U(IV), but U(V) is absent. In fact, significant amounts of structural Fe(II) are released from magnetite as Fe^{2+} ion to solution below $\text{pH} 6$,³⁰ leading to a decrease in the magnetite stoichiometry. Since non-stoichiometric magnetites are less efficient at incorporating metal ions (e.g. Ni(II) ³⁸, Co(II) ³⁹ and Cr(III) ³²), this process may explain the absence of U(V) stabilisation. At $\text{pH} 6$ (Figure 3e), the presence of U(IV) at intermediate stoichiometries, but its absence replaced by U(V) under the most reducing conditions ($R = 0.5$), supports this interpretation. At $\text{pH} 8$ (Figure 3f), strong interaction of U(IV) with magnetite, combined with unfavourable redox conditions for U(V) aqueous species, allows its occurrence. However, these mechanisms and conditions are not sufficient to overcome U(V) stabilisation, which occurs in similar proportions to U(IV). These results indicate that the stabilization of U(V) species on magnetite is strongly related to Fe(II) in magnetite structures, not only for the reduction process, but also for the U(V)-magnetite binding mechanism.

In a previous study involving conventional XANES, Latta et al.⁴ reported that reduction of U(VI) to UO_2 occurred only for $R > 0.38$ at $\text{pH} 7.2$. Our results show the presence of U(V) species on magnetite and that the U(VI) to U(V) reduction occurred at lower magnetite stoichiometry ($R > 0.2$). Besides the more difficult detection of U(V) by using normal XANES spectroscopy when U(VI) and U(IV) co-occur, this difference might be caused by the experimental conditions, such as the use of pH-buffer (50 mM MOPS or NaHCO_3) that might limit U(V) adsorption, and the higher U/Fe ratio (5 g L^{-1} magnetite and 500 μM U) that may favour the prevalence of U(IV) in the solid phase due to the enhanced precipitation of UO_2 . Our results suggest that the chemical species of U and its concentration in solution may also contribute to differences in the U adsorption mechanism on magnetite.

In conclusion, the present study provides new insights into interactions between U and magnetite nanoparticles regarding the influence of magnetite stoichiometry and pH on the U speciation. HERFD-XANES spectra of U(V) on magnetites show a peak splitting that has not been observed in conventional XANES, which enable us to demonstrate the high stability of U(V) species under a wide range of conditions, after 10 days on magnetite by the combination of MCR-ALS analysis. A key finding is that structural Fe(II), whose abundance depends on pH and redox conditions, plays a critical role in the stabilization of U(V) on magnetite. Compared with previous studies using conventional XANES, our results with the novel application of HERFD-XANES reveal that the reported U(VI) reduction of UO_2 only occurs for $R > 0.38$ at $\text{pH} 7.2$,⁴ highlighting the importance of high-resolution spectroscopy in accurately determining complex redox interactions at mineral surfaces. Revealing U(V) species, potentially acting a meta-stable species such as temporary sink or a reservoir for slow re-oxidation, on a widespread mineral like magnetite may be important to better predict the fate of U in reducing environments.

Data availability

The data supporting this article have been included in the IPGP Research Collection. See DOI: <https://doi.org/10.18715/IPGP.2025.mgb3600v>. Electronic Supplementary Information available: See DOI: 10.1039/x0xx00000x.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was also supported by the Grants-in-Aid for Scientific Research (KAKENHI) from JSPS (Grant Nos. 25K08534). The study was conducted with the approval of JASRI/SPring-8 (Proposal Nos. 2021B1821, 2022A1733 and 2024B1496). Additional support was provided by the COLOSSAL project funded by ANR (project number ANR-23-CE01-0001; PI: R.M.), as well as the French Brittany Region SAD program COLORED.

Notes and references

- P. L. Smedley and D. G. Kinniburgh, *Appl. Geochem.*, 2023, **148**, 105534.
- A. Y. Romanchuk, I. E. Vlasova and S. N. Kalmykov, *Front. Chem.*, 2020, **8**.
- Y. Takahashi, A. Yamaguchi and T. Yomogida, in *Treatise on Geochemistry (Third edition)*, eds. A. Anbar and D. Weis, Elsevier, Oxford, 2025, 105-150.
- D. E. Latta, C. A. Gorski, M. I. Boyanov, E. J. O'Loughlin, K. M. Kemner and M. M. Scherer, *Environ. Sci. Technol.*, 2012, **46**, 778-786.
- Z. Wang, K.-U. Ulrich, C. Pan and D. E. Giammar, *Environ. Sci. Technol. Lett.*, 2015, **2**, 227-232.
- H. E. Roberts, K. Morris, G. T. W. Law, J. F. W. Mosselmans, P. Bots, K. Kvashnina and S. Shaw, *Environ. Sci. Technol. Lett.*, 2017, **4**, 421-426.
- O. Stagg, K. Morris, A. Lam, A. Navrotsky, J. M. Velázquez, B. Schacherl, T. Vitova, J. Rothe, J. Galanzew, A. Neumann, P. Lythgoe, L. Abrahamsen-Mills and S. Shaw, *Environ. Sci. Technol.*, 2021, **55**, 16445-16454.
- S. Kushwaha, B. Sreedhar and P. Padmaja, *Langmuir*, 2012, **28**, 16038-16048.
- K. Yuan, E. S. Ilton, M. R. Antonio, Z. Li, P. J. Cook and U. Becker, *Environ. Sci. Technol.*, 2015, **49**, 6206-6213.
- C. Dewey, D. Sokaras, T. Kroll, J. R. Bargar and S. Fendorf, *Environ. Sci. Technol.*, 2020, **54**, 6021-6030.
- Z. Pan, B. Bartova, T. LaGrange, S. M. Butorin, N. C. Hyatt, M. C. Stennett, K. O. Kvashnina and R. Bernier-Latmani, *Nat. Commun.*, 2020, **11**, 4001.
- C. J. Dodge, A. J. Francis, J. B. Gillow, G. P. Halada, C. Eng and C. R. Clayton, *Environ. Sci. Technol.*, 2002, **36**, 3504-3511.
- K. Yuan, D. Renock, R. C. Ewing and U. Becker, *Geochim. Cosmochim. Acta*, 2015, **156**, 194-206.
- H. Veeramani, D. S. Alessi, E. I. Suvorova, J. S. Lezama-Pacheco, J. E. Stubbs, J. O. Sharp, U. Dippon, A. Kappler, J. R. Bargar and R. Bernier-Latmani, *Geochim. Cosmochim. Acta*, 2011, **75**, 2512-2528.
- Y. Wang, J. Wang, P. Li, H. Qin, J. Liang and Q. Fan, *Environ. Tech. Innov.*, 2021, **23**, 101615.
- R. Ding, C. Guida, C. I. Pearce, E. Arenholz, J.-M. Grenèche, A. Gloter, A. C. Scheinost, K. O. Kvashnina, K. Wang, A. Fernandez-Martinez, Y. Mu, K. M. Rosso and L. Charlet, *Sci. Adv.*, 2025, **11**, eadq3650.
- T. Vitova, K. O. Kvashnina, G. Nocton, G. Sukharina, M. A. Denecke, S. M. Butorin, M. Mazzanti, R. Caciuffo, A. Soldatov, T. Behrends and H. Geckeis, *Phys. Rev. B*, 2010, **82**, 235118.
- K. O. Kvashnina, S. M. Butorin, P. Martin and P. Glatzel, *Phys. Rev. Lett.*, 2013, **111**, 253002.
- K. O. Kvashnina and F. M. F. de Groot, *J. Electron Spectrosc. Relat. Phenom.*, 2014, **194**, 88-93.
- K. O. Kvashnina, Y. O. Kvashnin and S. M. Butorin, *J. Electron Spectrosc. Relat. Phenom.*, 2014, **194**, 27-36.
- J. G. Tobin, S. W. Yu, C. H. Booth, T. Tylliszczak, D. K. Shuh, G. van der Laan, D. Sokaras, D. Nordlund, T. C. Weng and P. S. Bagus, *Phys. Rev. B*, 2015, **92**, 035111.
- R. Bes, M. Rivenet, P. L. Solari, K. O. Kvashnina, A. C. Scheinost and P. M. Martin, *Inorg. Chem.*, 2016, **55**, 4260-4270.
- G. Leinders, R. Bes, J. Pakarinen, K. Kvashnina and M. Verwerft, *Inorg. Chem.*, 2017, **56**, 6784-6787.
- I. Pidchenko, K. O. Kvashnina, T. Yokosawa, N. Finck, S. Bahl, D. Schild, R. Polly, E. Bohnert, A. Rossberg, J. Gottlicher, K. Dardenne, J. Rothe, T. Schafer, H. Geckeis and T. Vitova, *Environ. Sci. Technol.*, 2017, **51**, 2217-2225.
- L. Desfougeres, E. Welcomme, M. Ollivier, P. M. Martin, J. Hennuyer, M. Hunault, R. Podor, N. Clavier and L. Favergeon, *Inorg. Chem.*, 2020, **59**, 8589-8602.
- R. Bes, G. Leinders and K. Kvashnina, *J. Synchrotron Radiat.*, 2022, **29**, 21-29.
- K. O. Kvashnina, Y. O. Kvashnin, J. R. Vegelius, A. Bosak, P. M. Martin and S. M. Butorin, *Anal. Chem.*, 2015, **87**, 8772-8780.
- T. Yomogida, D. Akiyama, K. Ouchi, Y. Kumagai, K. Higashi, Y. Kitatsuji, A. Kirishima, N. Kawamura and Y. Takahashi, *Inorg. Chem.*, 2022, **61**, 20206-20210.
- P. Jungcharoen, M. Pédrot, F. Choueikani, M. Pasturel, K. Hanna, F. Heberling, M. Tesfa and R. Marsac, *Environ. Sci.: Nano*, 2021, **8**, 2098-2107.
- P. Jungcharoen, M. Pédrot, F. Heberling, K. Hanna, F. Choueikani, C. Catrouillet, A. Dia and R. Marsac, *Environ. Sci.: Nano*, 2022, **9**, 2363-2371.
- P. Jungcharoen, R. Marsac, F. Choueikani, D. Masson and M. Pédrot, *Nanoscale Adv.*, 2023, **5**, 4213-4223.
- J. Scaria, M. Pédrot, L. Fablet, T. Yomogida, T. T. Nguyen, Y. Sivry, C. Catrouillet, A. E. Pradas Del Real, F. Choueikani, D. Vantelon, A. Dia, A. Groleau and R. Marsac, *Environ. Sci. Technol.*, 2025, **59**, 5747-5755.
- G. Landrot and E. Fonda, *J. Synchrotron Radiat.*, 2025, 1085-1094.
- B. Ravel and M. Newville, *J. Synchrotron Radiat.*, 2005, 537-541.
- W. Windig and D. A. Stephenson, *Anal. Chem.*, 1992, **64**, 2735-2742.
- H. Geckeis, J. Lützenkirchen, R. Polly, T. Rabung and M. Schmidt, *Chem. Rev.*, 2013, **113**, 1016-1062.
- R. Marsac, C. Catrouillet, M. Pédrot, M. F. Benedetti, A. Dia, E. D. van Hullebusch, M. Davranche, Y. Sivry, A.-C. Pierson-Wickmann, M. Tharaud and F. Heberling, *Current Opinion Colloid Interface Sci.*, 2024, **72**, 101820.
- L. Fablet, M. Pédrot, F. Choueikani, I. Kieffer, O. Proux, A.-C. Pierson-Wickmann, V. Cagniard, T. Yomogida and R. Marsac, *Environ. Sci.: Nano*, 2025, **12**, 2815-2827.
- L. Fablet, F. Choueikani, M. Pédrot and R. Marsac, *Environ. Sci.: Nano*, 2024, **11**, 2036-2048.

Data availability statements

The data supporting this article have been included in the IGP Research Collection See DOI: <https://doi.org/10.18715/IPGP.2025.mgb3600v>. Electronic Supplementary Information available: See DOI: 10.1039/x0xx00000x.